

A graph polynomial for independent sets of bipartite graphs

Qi Ge* Daniel Štefankovič*

February 10, 2010

Abstract

We introduce a new graph polynomial that encodes interesting properties of graphs, for example, the number of matchings and the number of perfect matchings. Most importantly, for bipartite graphs the polynomial encodes the number of independent sets ($\#BIS$).

We analyze the complexity of exact evaluation of the polynomial at rational points and show that for most points exact evaluation is $\#P$ -hard (assuming the generalized Riemann hypothesis) and for the rest of the points exact evaluation is trivial.

We conjecture that a natural Markov chain can be used to approximately evaluate the polynomial for a range of parameters. The conjecture, if true, would imply an approximate counting algorithm for $\#BIS$, a problem shown, by [10], to be complete (with respect to, so called, AP-reductions) for a rich logically defined sub-class of $\#P$. We give a mild support for our conjecture by proving that the Markov chain is rapidly mixing on trees. As a by-product we show that the “single bond flip” Markov chain for the random cluster model is rapidly mixing on constant tree-width graphs.

1 Introduction

Graph polynomials are a well-developed area useful for analyzing properties of graphs (see, e.g., the following survey papers [13, 14]). Arguably the most intriguing graph polynomial is the Tutte polynomial [38, 39]. The partition function of the random cluster model from statistical mechanics provides a particularly simple definition: for a graph $G = (V, E)$ let

$$Z(G; q, \mu) = \sum_{S \subseteq E} q^{\kappa(S)} \mu^{|S|}, \quad (1)$$

where $\kappa(S)$ is the number of connected components of the graph (V, S) . (It is well-known that the Tutte polynomial is obtained from Z by a simple transformation, see, e.g., equation (12) in Section 3.) The Tutte polynomial includes many graph polynomials as special cases, e.g., the chromatic polynomial, the flow polynomial, and the Potts model (see, e.g., [44]).

Now we define our graph polynomial.

Definition 1. *The R_2 -polynomial of a graph $G = (V, E)$ is*

$$R_2(G; q, \mu) = \sum_{S \subseteq E} q^{\text{rk}_2(S)} \mu^{|S|}, \quad (2)$$

*Department of Computer Science, University of Rochester, Rochester, NY 14627. Email: {qge, stefanko}@cs.rochester.edu. Research supported, in part, by NSF grant CCF-0910584.

where $\text{rk}_2(S)$ is the rank of the adjacency matrix of (V, S) over $\mathbb{F}_2$ (the field with 2 elements).

Now we look at how $R_2(G; q, \mu)$ encodes some properties of graphs. For $q = \mu^{-1/2}$ equation (2) becomes

$$P(G; \mu) := R_2(G; \mu^{-1/2}, \mu) = \sum_{S \subseteq E(G)} \mu^{|S| - \text{rk}_2(S)/2}.$$

We claim that $P(G; 0)$ is the number of matchings in G . To see this, note that $\text{rk}_2(S) \leq 2|S|$ (since adding an edge to S changes two entries in the adjacency matrix and hence can change rank by at most two), and $\text{rk}_2(S) < 2|S|$ if S is not a matching (since rank of the adjacency matrix of a star is $2 < 2|S|$, and adding further edges preserves the strict inequality).

Let

$$P(G; t, \mu) := t^{|V|} R_2(G; 1/t, \mu) \quad \text{and} \quad P_2(G; \mu) := \mu^{-|V|/2} P(G; 0, \mu).$$

Then $P_2(G; 0)$ is the number of perfect matchings of G . To see this, note that only subsets with full rank adjacency matrix contribute to $P(G; 0, \mu)$, and then only the minimal cardinality subsets with full rank adjacency matrix contribute to $P_2(G; 0)$ (these subsets are exactly the perfect matchings).

From now on we focus solely on bipartite graphs. For a *bipartite* graph $G = (U \cup W, E)$ we let

$$R'_2(G; \lambda, \mu) = \sum_{S \subseteq E} \lambda^{\text{rk}_2(S)} \mu^{|S|}, \quad (3)$$

where $\text{rk}_2(S)$ is the rank of the *bipartite* adjacency matrix of $(U \cup W, S)$. Note that

$$R_2(G; \lambda, \mu) = R'_2(G; \lambda^2, \mu), \quad (4)$$

since the adjacency matrix contains “two copies” of the bipartite adjacency matrix (one of them transposed). (The reason for definition (3) is that we prefer to operate with bipartite adjacency matrix for bipartite graphs.)

In Section 2 we prove that R'_2 counts the number of independent sets in bipartite graphs.

Theorem 1. *Let $G = (U \cup W, E)$ be a bipartite graph. The number of independent sets of G is given by*

$$2^{|U|+|W|-|E|} R'_2(G; 1/2, 1).$$

We show in Section 3 that exact evaluation of the polynomial $R'_2(G; \lambda, \mu)$ is #P-hard at a variety of rational points (λ, μ) assuming the validity of the generalized Riemann hypothesis (GRH).

Theorem 2. *Exact evaluation of R'_2 at rational point (λ, μ) is*

- *polynomial-time computable when $\lambda \in \{0, 1\}$ or $\mu = 0$ or $(\lambda, \mu) = (1/2, -1)$;*
- *#P-hard when $\lambda \notin \{0, 1, 1/2\}$ and $\mu \neq 0$, assuming GRH;*
- *#P-hard when $\lambda = 1/2$ and $\mu \notin \{0, -1\}$.*

Remark 1. *For the non-bipartite case we have the following classification. Exact evaluation of R_2 at rational point (λ, μ) is polynomial-time computable when $\mu = 0$ or $\lambda \in \{-1, 0, 1\}$; the $\lambda = -1$ case follows from the fact that a skew-symmetric matrix with zero diagonal has even rank over any field (the zero diagonal condition is redundant for fields of characteristic $\neq 2$). For any other*

rational λ and μ we get #P-hardness of evaluating the R_2 polynomial from Theorem 2 and (4) (again assuming GRH). (Note that $(\lambda, \mu) \mapsto (\lambda^2, \mu)$ never maps to the easy case $(1/2, -1)$, since λ is rational. It would be nice to have hardness classification of evaluating R_2 and R'_2 for, say, algebraic λ and μ .)

Because of the hardness of exact evaluation of R'_2 , we turn to approximate evaluation of $R'_2(G; \lambda, \mu)$.

Remark 2. *There is a fully polynomial randomized approximation scheme (FPRAS) for $R'_2(G; \lambda, \mu)$ when $\lambda = 1/2$ and $-1 < \mu < 0$. This follows from Theorem 5 (stated in Section 3) and the fact that there is an FPRAS for #PBIS(η) (defined in Section 3) when $0 < \eta < 1$ (see [19], and note that #PBIS(η) corresponds to $\beta = 1$, $\gamma = (1 + \eta)/(1 - \eta)$ and $\mu = 1$ in their parametrization).*

We now define the sampling problem associated with R'_2 .

RANK WEIGHED SUBGRAPHS with $\lambda, \mu \geq 0$, (RWS(λ, μ))

Instance: a bipartite graph $G = (U \cup W, E)$,

Output: $S \subseteq E$ with probability of $S \propto \lambda^{\text{rk}_2(S)} \mu^{|S|}$.

The “single bond flip” chain is a natural approach to sampling from RWS(λ, μ).

Definition 2. *Single bond flip chain is defined as follows: pick an edge $e \in E$ at random and let $S = X_t \oplus \{e\}$. Set $X_{t+1} = S$ with probability $(1/2) \min\{1, \lambda^{\text{rk}_2(S) - \text{rk}_2(X_t)} \mu^{|S| - |X_t|}\}$ and $X_{t+1} = X_t$ with the remaining probability.*

In each step of the single bond flip chain, we have to compute the rank of a matrix over $\mathbb{F}_2$ (corresponding to S) which differs from the current matrix (corresponding to X_t) in a single entry. One can use dynamic matrix rank problem algorithms to perform this computation in $O(n^{1.575})$ arithmetic operations per step [16].

Instead of flipping one edge in a step, we can have another Markov chain which flips a random subset of edges adjacent to a single vertex. It seems likely that the new chain can generate good random samples faster than the single bond flip chain—a step of the new chain can be performed in $O(n^2)$ arithmetic operations (using “rank one update” for the dynamic matrix rank problem [16]).

We optimistically conjecture that the single bond flip chain (and hence the other chain) mixes rapidly.

Conjecture 1. *For any fixed $\lambda, \mu > 0$ the single bond flip chain mixes in polynomial time.*

Note that Goldberg and Jerrum [18] conjecture the opposite, in particular for $\lambda = 1/2$ and $\mu = 1$. We make the conjecture based on similarity of R'_2 polynomial to the random cluster model and the fact that slow mixing of the single bond flip chain was not established in random cluster model (slow mixing is usually easier to establish than rapid mixing).

Conjecture 1 could be false for small λ and large μ if the problem of maximizing $|S| - c \cdot \text{rk}_2(S)$ (c is a constant and $c > 1$) over $S \subseteq E$ is hard (the corresponding maximization problem is easy for the random cluster model).

We can provide a mild support for Conjecture 1. In Section 5 we prove that for fixed $\lambda, \mu > 0$ the single bond flip chain mixes, in time polynomial in the number of vertices, for trees.

Theorem 3. *For every fixed $\lambda, \mu > 0$, the mixing time $\tau(\varepsilon)$ of the single bond flip chain for a tree on n vertices is*

$$\tau(\varepsilon) = O\left(n^{3 + \lceil \log_2 \lambda \rceil} (|\log \lambda| + |\log \mu| + \log(1/\varepsilon))\right).$$

As a by-product of our techniques, we show that single bond flip Markov chain for the random cluster model is rapidly mixing if $q, \mu > 0$ and G has constant tree-width (the condition $q, \mu > 0$ is equivalent to $x, y > 1$ for the Tutte polynomial $T(G; x, y)$).

2 Independent sets in bipartite graphs

The problem of counting independent sets ($\#IS$) in a graph is of interest in both computer science and statistical physics (independent sets are a special case of the so-called hard-core model, see, e.g., [5]). Exact computation of $\#IS$ is $\#P$ -complete even for 3-regular planar bipartite graphs [40, 45]. Fully polynomial randomized approximation scheme (FPRAS) is known for graphs with maximum degree $\Delta \leq 5$, [28, 12, 43]. Unless $RP=NP$, an FPRAS does not exist for graphs with $\Delta \geq 25$, [9].

Now we focus on the problem of counting independent sets in bipartite graphs ($\#BIS$). While for exact counting the complexity of $\#BIS$ and $\#IS$ is the same, the situation looks very different for approximate counting, for example, no inapproximability result is known for $\#BIS$. Dyer et al. [10] show that $\#BIS$ is complete w.r.t. approximation-preserving reductions (AP-reductions) in a sub-class of $\#P$. Many problems were shown to be equivalent (w.r.t. AP-reductions) to $\#BIS$, for example, $\#DOWNSETS$, $\#1P1NSAT$ [10], computing the partition function of a ferromagnetic Ising model with local fields [17], and counting the number of satisfying assignments of a class of Boolean CSP instances [11]. A pertinent negative result for $\#BIS$ is that Glauber dynamics (or more generally, any chain whose states are independent sets and that flips at most $0.35n$ vertices in one step) cannot be used to efficiently sample random independent sets in a random 6-regular bipartite graphs on $n + n$ vertices [9].

The rest of this section is devoted to proving Theorem 1. It will be convenient to work with matrices instead of graphs. For two zero-one matrices A, B we say $B \leq A$ if B corresponds to a subgraph of A , formally

Definition 3. Let A, B be zero-one $n_1 \times n_2$ matrices. We say $B \leq A$ if $A_{ij} = 0$ implies $B_{ij} = 0$, for all $i \in [n_1]$ and $j \in [n_2]$. Let $\mathcal{C}_A$ be the set of zero-one $n_1 \times n_2$ matrices B such that $B \leq A$.

Let $\#_1(A)$ denote the number of ones in A (that is, the number of edges in the corresponding graph). The RWS problem rephrased for matrices is:

RANK WEIGHED MATRICES with $\lambda, \mu \geq 0$ ($RWM(\lambda, \mu)$)

Instance: an $n_1 \times n_2$ matrix A .

Output: $B \in \mathcal{C}_A$ with probability of $B \propto \lambda^{\text{rk}_2(B)} \mu^{\#_1(B)}$.

The problem of sampling independent sets in bipartite graphs is:

BIPARTITE INDEPENDENT SETS (BIS)

Instance: a bipartite graph $G = (U \cup W, E)$.

Output: a uniformly random independent set of G .

Before we show a connection between BIS and $RWM(1/2, 1)$ we remark that to sample bipartite independent sets it is enough to sample a subset of one side, say U , from the correct (marginal) distribution. We now describe this distribution in a setting which will be advantageous for the proof of Theorem 1.

We will represent an independent set by a pair of (indicator) vectors u, v (where $u \in \mathbb{F}_2^{n_1}$ and $v \in \mathbb{F}_2^{n_2}$).

Definition 4. We say that two vectors $\alpha, \beta \in \mathbb{F}_2^n$ share a one if there exists $i \in [n]$ such that $\alpha_i = \beta_i = 1$.

We will use the following simple fact.

Observation 1. Let $\alpha, \beta \in \mathbb{F}_2^n$. Let d be the number of ones in β .

- if α, β share a one then there are 2^{d-1} vectors $\beta' \leq \beta$ such that $\alpha^T \beta' \equiv 0 \pmod{2}$.
- if α, β do not share a one then there are 2^d vectors $\beta' \leq \beta$ such that $\alpha^T \beta' \equiv 0 \pmod{2}$.

Let $u \in \mathbb{F}_2^{n_1}$ be a vector. We would like to count the number of $v \in \mathbb{F}_2^{n_2}$ such that u, v is an independent set. Note that u, v is an independent set iff $v_j = 0$ for every $j \in [n_2]$ such that u and j -th column of A share a one. Let k be the number of columns of A that do not share a one with u . Then we have

$$u \in \mathbb{F}_2^{n_1} \text{ occurs in } 2^k \text{ independent sets.} \quad (5)$$

Thus to sample independent sets in a bipartite graph G with $n_1 \times n_2$ bipartite adjacency matrix A it is enough to sample $u \in \mathbb{F}_2^{n_1}$ with the probability of u proportional to 2^k , where k is the number of columns of A that do not share a one with u . We will call this distribution on u the marginal BIS distribution.

The following lemma shows a tight connection between BIS and RWM(1/2, 1)—given a sample from one distribution it is trivial to obtain a sample from the other one.

Lemma 1. Let G be a bipartite graph with bipartite adjacency matrix A .

- Let u, v be a uniformly random independent set of G . Let B be a uniformly random matrix from the following set $\{D \in \mathcal{C}_A \mid u^T D \equiv 0 \pmod{2}\}$. Then B is from the RWM(1/2, 1)-distribution.
- Let $B \in \mathcal{C}_A$ be a random matrix from the RWM(1/2, 1)-distribution. Let $u \in \mathbb{F}_2^{n_1}$ be a uniformly random vector from the left null space of B (that is, $\{\beta \in \mathbb{F}_2^{n_1} \mid \beta^T B \equiv 0 \pmod{2}\}$). Then u is from the marginal BIS distribution.

Proof. Let Q be the set of u, B pairs such that $u^T B \equiv 0 \pmod{2}$ and $B \leq A$. Let ψ be the uniform distribution on Q . Note that ψ marginalized over u yields the RWM(1/2, 1)-distribution on $B \leq A$, here we are using the fact that a d -dimensional space (in this case the left null space of B) over $\mathbb{F}_2$ has 2^d elements. Formally,

$$P(B) = \sum_{u: u^T B \equiv 0 \pmod{2}} \frac{1}{|Q|} = \frac{2^{n_1 - \text{rk}_2(B)}}{|Q|} = \frac{2^{-\text{rk}_2(B)}}{R'_2(G; 1/2, 1)}. \quad (6)$$

Next we show that ψ marginalized over B yields the marginal BIS distribution. We compute the number of $B \leq A$ such that $u^T B \equiv 0 \pmod{2}$. Let us use the same k as in (5), that is, k is the number of columns of A that do not share a one with u .

Note that the columns of B can be chosen independently and only if the column and u share a one is the number of choices (for that column) halved. Let $\#_1(A)$ be the number of ones in A . Thus

$$\text{there are } 2^{\#_1(A) - (n_2 - k)} \text{ choices of } B \leq A \text{ such that } u^T B \equiv 0 \pmod{2}. \quad (7)$$

Note that for fixed u the counts in (5) and (7) differ by a factor of $2^{\#_1(A)-n_2}$ (which is independent of u). Thus ψ marginalized over B yields the marginal BIS distribution on u . Formally

$$P(u) = \frac{2^{\#_1(A)-(n_2-k)}}{|Q|} = \frac{2^k}{\#\text{BIS}(G)}. \quad (8)$$

Note that this proves both claims of the lemma since in both cases the u, B pair is from ψ (by first sampling from a marginal and then sampling the remaining variable) and the conclusion in both claims is a statement about marginal (of the remaining variable). $\square$

Theorem 1 now follows from the proof of Lemma 1.

Proof of Theorem 1. Let Q be the set from the proof of Lemma 1. From (6) we obtain

$$|Q| = R'_2(G; 1/2, 1)2^{n_1}. \quad (9)$$

From (8) we have that the number of independent sets of G is given by

$$\#\text{BIS}(G) = \frac{|Q|}{2^{\#_1(A)-n_2}}. \quad (10)$$

Combining (9) and (10) we obtain the theorem. $\square$

We do not know a good combinatorial interpretation for the mod-2 rank of B for general graphs. For forests (which are, of course, always bipartite) we have the following characterization.

Lemma 2. *Let $G = (V, E) = (U \cup W, E)$ be a forest with bipartite adjacency matrix A . Then $\text{rk}_2(A)$ is the size of maximum matching in G .*

Proof. Let $a \in V$ be a leaf of G and let $e = \{a, b\} \in E$ be the edge adjacent to a . Note that b is matched in every maximum matching M (otherwise one could add e to M). Thus removing b and all adjacent edges decreases the size of maximum matching by 1.

Now we argue that removing b (and all adjacent edges) also decreases rank (over $\mathbb{F}_2$) by 1. W.l.o.g. assume that b corresponds to the first row and a corresponds to the first column. Removing b (and all adjacent edges) corresponds to removing the first row of A . Note that this decreases rank by at most 1 and it does decrease it by 1, since the only non-zero entry in the first column is in the first row. $\square$

3 Exact evaluation of R'_2 (proof of Theorem 2)

We will prove Theorem 2 in this section. Let $G = (V, E) = (U \cup W, E)$ be a bipartite graph. First we deal with the cases where exact evaluation of $R'_2(G; \lambda, \mu)$ is easy. For cases $\lambda \in \{0, 1\}$ and $\mu = 0$ we have

$$R'_2(G; 0, \mu) = R'_2(G; \lambda, 0) = 1 \quad \text{and} \quad R'_2(G; 1, \mu) = (1 + \mu)^{|E|}.$$

For $\lambda = 1/2$ and $\mu = -1$ we will show

$$R'_2(G; 1/2, -1) = 2^{|E|-|V|+t}, \quad (11)$$

where t is the number of isolated vertices in G (see the remark after Theorem 5 in this section).

For the hardness results we give reductions from the problem of evaluating the Tutte polynomial (to establish the second part of Theorem 2) and #BIS (to establish the third part of Theorem 2).

The Tutte polynomial of a graph $G = (V, E)$ is a polynomial in two variables x, y defined by

$$T(G; x, y) = \sum_{S \subseteq E} (x-1)^{\kappa(S)-\kappa(E)} (y-1)^{|S|-|V|+\kappa(S)},$$

where $\kappa(S)$ is the number of connected components of the graph (V, S) . The Tutte polynomial is closely related to the random cluster model (see, e.g., [44]). Let $Z(G; q, \mu)$ be defined as in (1). One has

$$T(G; x, y) = (x-1)^{-\kappa(E)} (y-1)^{-|V|} Z(G; (x-1)(y-1), (y-1)), \quad (12)$$

where we assume $x \neq 1$ and $y \neq 1$.

We are going to use the following result on the complexity of exact evaluation of the Tutte polynomial.

Theorem 4 ([23]). *Exact evaluation of the Tutte polynomial is #P-hard for all rational numbers x, y except when*

1. $(x-1)(y-1) = 1$; or
2. (x, y) equals $(1, 1)$, $(-1, -1)$, $(0, -1)$ or $(-1, 0)$.

The second part of Theorem 2 will be proved by reducing from exact evaluation of the Tutte polynomial. We prove the following Lemma in Section 3.1.

Lemma 3. *Assuming the validity of the GRH, exact evaluation of the Tutte polynomial at x, y is polynomial-time Turing reducible to exact evaluation of R'_2 at λ, μ , when*

$$(x-1)(y-1) = 1/\lambda - 1, \quad y-1 = \mu^2, \quad \lambda \notin \{0, 1\}, \quad \text{and} \quad \mu \neq 0. \quad (13)$$

Assuming GRH, by Lemma 3 and Theorem 4, we have that exact evaluation of R'_2 at rational point (λ, μ) is #P-hard when $\lambda \notin \{0, 1/2, 1\}$ and $\mu \neq 0$. We do not get #P-hardness for $\lambda = 1/2$ since the reduction is from the Tutte polynomial at $(x-1)(y-1) = 1$ which is polynomial-time computable (part 1 of Theorem 4). (The other easy cases of the Tutte polynomial have no impact since $y = 1$ implies $\mu = 0$ and $y \in \{0, -1\}$ implies that μ is not real.) We proved the second part of Theorem 2.

Now we prove the third part of Theorem 2 (the proof of main lemmas is deferred to later sections). To show #P-hardness of exact evaluation of $R'_2(G; 1/2, \mu)$ for $\mu \notin \{-1, 0\}$, we prove a connection between R'_2 and the “permissive version of #BIS” (#PBIS) introduced in [17]; #PBIS is a generalization of #BIS where the weight of a set of vertices is determined by the number of pairs of neighboring vertices that are both in the set (in #BIS the weight is zero raised to the number of such pairs).

#PERMISSIVE BIPARTITE INDEPENDENT SETS with η (#PBIS(η))

Instance: a bipartite graph $G = (U \cup W, E)$.

Output: the quantity

$$\#PBIS(G; \eta) = \sum_{\sigma: U \cup W \rightarrow \{0, 1\}} (1 + \eta)^{w(\sigma)} (1 - \eta)^{|E| - w(\sigma)},$$

where $w(\sigma)$ is the number of edges in E with both endpoints labelled 1 by σ .

(We are using a different parametrization than [17]—our η and their γ are connected by $\gamma^2 = (1 + \eta)/(1 - \eta)$.)

Note that

$$\#\text{BIS}(G) = 2^{|E|} \#\text{PBIS}(G; -1).$$

The following result is a generalization of Theorem 1 and shows that R'_2 encodes $\#\text{PBIS}(\eta)$ as well. The proof is deferred to Section 3.2.

Theorem 5. *Let $G = (V, E) = (U \cup W, E)$ be a bipartite graph.*

$$\#\text{PBIS}(G; \eta) = 2^{|V|} R'_2(G; 1/2, -\eta).$$

Note that $\#\text{PBIS}(G; 1) = 2^{|E|+t}$, where t is the number of isolated vertices of G (since the other vertices have to be labeled 1 by σ). This implies (11).

The following result on $\#\text{PBIS}(\eta)$ will be proved in Section 3.3.

Lemma 4. *$\#\text{BIS}$ is polynomial-time Turing reducible to $\#\text{PBIS}(\eta)$ with η a rational number and $\eta \notin \{\pm 1, 0\}$.*

The third part of Theorem 2 follows from Theorem 5, Lemma 4 and the fact that exact computation of $\#\text{BIS}$ is $\#\text{P}$ -complete [33].

3.1 Reducing Tutte polynomial to R'_2 polynomial (proof of Lemma 3)

We will focus on bipartite graphs $G = (U \cup W, E)$ such that vertices in partition W have degree at most 2 (a natural operation that produces such graphs is *2-stretch*, that is, replacement of each edge with a path of length 2).

Let $G = (U \cup W, E)$ be a bipartite graph with max-degree in W bounded by 2. We call a connected component $C = (U_C \cup W_C, E_C)$ of G *pure* if every vertex in W_C has degree 2 in C . A component that is not pure will be called *mixed*. The evaluation of R'_2 polynomial in G can be expressed using pure connected components as follows.

Lemma 5. *For every bipartite graph $G = (U \cup W, E)$ such that the degree of each vertex in W is bounded by 2,*

$$R'_2(G; \lambda, \mu) = \sum_{S \subseteq E} \lambda^{|U| - \kappa'(S)} \mu^{|S|},$$

where $\kappa'(S)$ is the number of pure connected components in $(U \cup W, S)$.

Before proving Lemma 5 we need the following characterization of the rank of bipartite adjacency matrices over $\mathbb{F}_2$.

Lemma 6. *Let $G = (U \cup W, E)$ be a connected bipartite graph with max-degree in W bounded by 2. Let B be the adjacency matrix of G . Then*

$$\text{rk}_2(B) = \begin{cases} |U| & \text{if there is a vertex of degree 1 in } W, \\ |U| - 1 & \text{otherwise.} \end{cases}$$

Proof. Let $x \in \mathbb{F}_2^U$ be a solution of the linear system $x^T B = 0$. Let U_i be the set of vertices $u \in U$ such that $x_u = i$, for $i = 0, 1$. Note that no vertex $v \in W$ has neighbors both in U_0 and U_1 (otherwise $(x^T B)_v = 1$). Thus for G to be connected either $x = 0$ or $x = 1$. If there is a vertex of degree 1 in W then $x = 1$ is not a solution and hence $\text{rk}_2(B) = |U|$. On the other hand if all vertices in W have degree 2 then $x = 1$ is a solution and hence $\text{rk}_2(B) = |U| - 1$. $\square$

Now we prove Lemma 5.

Proof of Lemma 5. We will show that

$$\text{rk}_2(S) = |U| - \kappa'(S), \quad (14)$$

where $\text{rk}_2(S)$ is the rank (over $\mathbb{F}_2$) of B , the bipartite adjacency matrix of $(U \cup W, S)$, and $\kappa'(S)$ is the number of pure connected components of $(U \cup W, S)$.

Note that B has a block structure with a block for each connected component. The rank is the sum of the ranks of the blocks. Equation (14) now follows from Lemma 6. $\square$

We now lay groundwork for the proof of Lemma 3. We use the following construction in the reduction. Given a graph $H = (V_H, E_H)$ and a bipartite graph $\Upsilon = (U_\Upsilon \cup W_\Upsilon, E_\Upsilon)$ with a specific vertex $u \in U_\Upsilon$, we construct a bipartite graph G from H and Υ as follows. Let $\hat{H} = (U_{\hat{H}} \cup W_{\hat{H}}, E_{\hat{H}})$ be the 2-stretch of H , where $U_{\hat{H}}$ corresponds to the vertices of H . For each vertex v in $U_{\hat{H}}$ we identify v with u in a copy of Υ (thus we have $|V_H| = |U_{\hat{H}}|$ copies of Υ). We call the graph G the *stretch-sum* of H and (Υ, u) . Note that if W_Υ contains only vertices of degree at most 2 then the partition of G containing $W_{\hat{H}}$ contains only vertices of degree at most 2.

We define two functions related to R'_2 .

Definition 5. Let $\lambda, \mu \in \mathbb{R}$. Let $\Upsilon = (U \cup W, E)$ be a bipartite graph with a specific vertex $u \in U$. Assume that the max-degree in W bounded by 2. We define

$$Z'_p(\Upsilon; \lambda, \mu) = \sum_S \lambda^{-\kappa'(S)} \mu^{|S|}, \quad (15)$$

where the sum is over all $S \subseteq E$ such that u is in a pure connected component of $(U \cup W, S)$, and $\kappa'(S)$ is the number of pure connected components of $(U \cup W, S)$.

Similarly, we define

$$Z'_m(\Upsilon; \lambda, \mu) = \sum_S \lambda^{-\kappa'(S)} \mu^{|S|}, \quad (16)$$

where the sum is over all sets $S \subseteq E$ such that u is in a mixed connected component of $(U \cup W, S)$.

The following lemma provides a connection between the random cluster partition function Z of H and the R'_2 polynomial of G for rational λ and μ .

Lemma 7. Fix rational $\lambda \notin \{0, 1\}$ and rational $\mu \neq 0$. Let p be a prime such that $\lambda \in \mathbb{Z}_p^*$. Let $\Upsilon = (U \cup W, E)$ be a bipartite graph with a specific vertex $u \in U$, such that the max-degree in W bounded by 2. Suppose Υ satisfies

$$\begin{aligned} X &:= \lambda Z'_p(\Upsilon; \lambda, \mu) \not\equiv 0 \pmod{p}, \\ Y &:= Z'_m(\Upsilon; \lambda, \mu) + \lambda Z'_p(\Upsilon; \lambda, \mu) \equiv 0 \pmod{p}. \end{aligned} \quad (17)$$

Let G be the stretch-sum of $H = (V_H, E_H)$ and (Υ, u) . Then

$$R'_2(G; \lambda, \mu) \equiv \lambda^{|V_H| \cdot |U|} X^{|V_H|} Z(H; 1/\lambda - 1, \mu^2) \pmod{p}, \quad (18)$$

where $Z(H; 1/\lambda - 1, \mu^2)$ is defined in (1).

Proof. Let $\hat{H} = (V_{\hat{H}}, E_{\hat{H}})$ be the 2-stretch of H . Note that $\hat{H}$ is a subgraph of $G = (V_G, E_G)$. Let $S_0 \subseteq E_{\hat{H}}$. Let Λ_{S_0} be a family of subsets $S \subseteq E_G$ such that $S \cap E_{\hat{H}} = S_0$. Now, we evaluate

$$\sum_{S \in \Lambda_{S_0}} \lambda^{-\kappa'(S)} \mu^{|S|} \quad (19)$$

modulo p .

Claim 1.

$$(19) \equiv \begin{cases} (\lambda^{-1} - 1)^{\kappa(S_0)} \mu^{|S_0|} X^{|V_H|} & \text{mod } p \quad \text{if } (V_{\hat{H}}, S_0) \text{ has no mixed connected component,} \\ 0 & \text{mod } p \quad \text{otherwise,} \end{cases}$$

where $\kappa(S_0)$ is the number of connected components of $(V_{\hat{H}}, S_0)$.

Proof of Claim 1. Equation (19) can be rewritten as a product, where each term in the product corresponds to a connected component of $(V_{\hat{H}}, S_0)$ (since each connected component with the copies of Υ attached to it influences $\kappa'(S)$ independently). Thus

$$(19) = \prod_C \Phi_C,$$

where for each connected component $C = (V_C, E_C)$ in $(V_{\hat{H}}, S_0)$ such that there are k copies of Υ (we refer to the copies $\Upsilon_1, \dots, \Upsilon_k$ and to their special vertices $u_1, \dots, u_k$) attached to it,

$$\Phi_C = \sum_{S_1 \subseteq E_{\Upsilon_1}} \dots \sum_{S_k \subseteq E_{\Upsilon_k}} \lambda^{-\kappa'(\bigcup_{i \in [k]} S_i \cup E_C)} \mu^{|E_C| + \sum_{i \in [k]} |S_i|}. \quad (20)$$

Let $A_{i,0}$ be the set of S_i such that u_i is in a mixed component of (V_{Υ_i}, S_i) and let $A_{i,1} = 2^{E_{\Upsilon_i}} \setminus A_{i,0}$. Equation (20) can be written as follows

$$\Phi_C = \sum_{x_1=0}^1 \dots \sum_{x_k=0}^1 \sum_{S_1 \in A_{1,x_1}} \dots \sum_{S_k \in A_{k,x_k}} \lambda^{-\kappa'(\bigcup_{i \in [k]} S_i \cup E_C)} \mu^{|E_C| + \sum_{i \in [k]} |S_i|}. \quad (21)$$

We have

$$\kappa' \left(\bigcup_{i \in [k]} S_i \cup E_C \right) = \sum_{i \in [k]} \kappa'_i(S_i) - \sum_{i=1}^k x_i + \ell,$$

where $\ell = 1$ if $x_1 = \dots = x_k = 1$ and $C = (V_C, E_C)$ is a pure connected component of $(V_{\hat{H}}, S_0)$, and $\ell = 0$ otherwise. Thus

$$\begin{aligned} & \sum_{S_1 \in A_{1,x_1}} \dots \sum_{S_k \in A_{k,x_k}} \lambda^{-\kappa'(\bigcup_{i \in [k]} S_i \cup E_C)} \mu^{|E_C| + \sum_{i \in [k]} |S_i|} = \\ & \lambda^{-\ell} \mu^{|E_C|} \sum_{S_1 \in A_{1,x_1}} \dots \sum_{S_k \in A_{k,x_k}} \prod_{i=1}^k \lambda^{-\kappa'_i(S_i) + x_i} \mu^{|S_i|} = \lambda^{-\ell} \mu^{|E_C|} X^{k'} (Y - X)^{k-k'}, \end{aligned} \quad (22)$$

where $k' = x_1 + \dots + x_k$.

Plugging (22) into (21) we obtain

$$\Phi_C = \mu^{|E_C|} Y^k + L, \quad (23)$$

where $L = (1/\lambda - 1)\mu^{|E_C|} X^k$ if C is a pure component of $(V_{\hat{H}}, S_0)$ and 0 otherwise. Evaluating (23) modulo p (using (17)) we obtain

$$\Phi_C \equiv \begin{cases} 0 \pmod{p} & \text{if } C \text{ is a mixed component of } (V_{\hat{H}}, S_0), \\ (1/\lambda - 1)\mu^{|E_C|} X^k \pmod{p} & \text{otherwise.} \end{cases}$$

Thus (19) is zero modulo p if there is a mixed component C in $(V_{\hat{H}}, S_0)$. Assume now that all components of $(V_{\hat{H}}, S_0)$ are pure. The total number of edges in the components is $|S_0|$, the total number of copies of Υ in the components is $|V_H|$, and hence

$$(19) \equiv (1/\lambda - 1)^{\kappa(S_0)} \mu^{|S_0|} X^{|V_H|} \pmod{p}.$$

□

Now we use Claim 1 to prove (18). Note that by Lemma 5,

$$R'_2(G; \lambda, \mu) = \lambda^{|V_H| \cdot |U|} \sum_{S_0 \subseteq E_{\hat{H}}} \sum_{S \in \Lambda_{S_0}} \lambda^{-\kappa'(S)} \mu^{|S|}, \quad (24)$$

where $\kappa'(S)$ is the number of pure connected components of (V_G, S) .

Note that by Claim 1 if S_0 contains a mixed component then the inner sum in (24) is 0 modulo p . Thus to evaluate (24) modulo p it is enough to sum over S_0 which contain only pure components. Each such S_0 is obtained from exactly one $S' \subseteq E_H$ by 2-stretching. Note $(V_{\hat{H}}, S_0)$ has the same number of connected components as (V_H, S') , and $|S_0| = 2|S'|$. Thus

$$\begin{aligned} R'_2(G; \lambda, \mu) &\equiv \lambda^{|V_H| \cdot |U|} X^{|V_H|} \sum_{S' \subseteq E_H} (1/\lambda - 1)^{\kappa(S')} \mu^{2|S'|} \\ &\equiv \lambda^{|V_H| \cdot |U|} X^{|V_H|} Z(H; 1/\lambda - 1, \mu^2) \pmod{p}. \end{aligned}$$

□

We use different Υ for different values of μ in the reduction. When $\mu \neq -2$ we let Υ_1 be the bipartite graph with bipartition $U = \{u_0, u_1\}$, $W = \{v_i \mid 0 \leq i \leq k\}$ and $k+2$ edges: edge $\{u_0, v_0\}$, and an edge between u_1 and each v_i , for $0 \leq i \leq k$. The specific vertex of Υ_1 is u_0 . By elementary counting, we have

$$\begin{aligned} \lambda Z'_p(\Upsilon_1; \lambda, \mu) &= (\mu + 1)^{k+1} + \mu^2 + \lambda^{-1} - 1, \\ \lambda Z'_p(\Upsilon_1; \lambda, \mu) + Z'_m(\Upsilon_1; \lambda, \mu) &= (\mu + 1)((\mu + 1)^{k+1} + \lambda^{-1} - 1). \end{aligned} \quad (25)$$

When $\mu = -2$ we let Υ_2 be the bipartite graph with bipartition $U = \{u_0, u_1, u_2\}$, $W = \{v_i \mid 0 \leq i \leq 2k\}$ and $4k+2$ edges: $\{u_0, v_0\}$, $\{u_1, v_0\}$, and a complete bipartite graph between $U \setminus \{u_0\}$ and $W \setminus \{v_0\}$. The specific vertex of Υ_2 is u_0 . By elementary counting, we have

$$\begin{aligned} \lambda Z'_p(\Upsilon_2; \lambda, -2) &= \lambda^{-2} + 5^{2k} \lambda^{-1} - 3 + 3 \cdot 5^{2k} + \lambda^{-1}, \\ \lambda Z'_p(\Upsilon_2; \lambda, -2) + Z'_m(\Upsilon_2; \lambda, -2) &= -\lambda^{-2} - 5^{2k} \lambda^{-1} - 1 + 5^{2k} + 3\lambda^{-1}. \end{aligned} \quad (26)$$

Fix rational $\lambda \notin \{0, 1\}$ and $\mu \neq 0$. We want to find sufficiently many primes such that there is some integer k for which (25) satisfies (17) (when $\mu \neq -2$) or (26) satisfies (17) (when $\mu = -2$). We need the following result on the density of primes.

Lemma 8 ([29, 30, 35]). *Let $r, q \in \mathbb{Q}^*$ and $q \neq \pm 1$. The density (inside the set of all primes) of primes p such that*

$$q^k \equiv r \pmod{p},$$

can be satisfied for some integer k is a positive constant, assuming GRH.

Lemma 8 immediately yields the following two corollaries.

Corollary 1. *Fix rational $\lambda \notin \{0, 1\}$ and rational $\mu \notin \{0, -2\}$. The density (inside the set of all primes) of the primes p such that there is an integer k for which (25) satisfies (17) is a positive constant, assuming GRH.*

Proof. If $\mu = -1$, then to make (25) satisfy (17), it is sufficient to have $\lambda^{-1} \not\equiv 0 \pmod{p}$. Thus, for all but a constant numbers of primes, and for all positive integers k , (25) satisfies (17).

Now assume $\mu \neq -1$. To make (25) satisfy (17), it is sufficient to have

$$\begin{aligned} (\mu + 1)^{k+1} + \lambda^{-1} - 1 &\equiv 0 \pmod{p}, \\ \mu &\not\equiv 0 \pmod{p}. \end{aligned} \tag{27}$$

The corollary follows from Lemma 8 (and the fact that the number of primes such that $\mu \equiv 0 \pmod{p}$ is finite). $\square$

Corollary 2. *Fix rational $\lambda \notin \{0, 1\}$ and rational $\mu = -2$. The density (inside the set of all primes) of the primes p such that there is an integer k for which (26) satisfies (17) is a positive constant, assuming GRH.*

Proof. We claim that if

$$\begin{aligned} (\lambda^{-2} - 3\lambda^{-1} + 1)(1 - \lambda^{-1})^{-1} &\equiv 25^k \pmod{p}, \\ 1 - \lambda^{-1} &\not\equiv 25^k \pmod{p} \end{aligned} \tag{28}$$

then (26) satisfies (17). The first equation in (28) makes the second equation in (17) satisfied; and the first and the second equation in (28) make the first equation in (17) satisfied.

By Lemma 8 the density of primes that make the first equation in (28) satisfied is positive. Solving

$$(\lambda^{-2} - 3\lambda^{-1} + 1)(1 - \lambda^{-1})^{-1} \equiv 1 - \lambda^{-1} \pmod{p},$$

we obtain $\lambda^{-1} \equiv 0 \pmod{p}$ and hence the second equation in (28) is automatically satisfied. $\square$

Proof of Lemma 3. Let x, y, λ, μ be rational numbers such that (13) is satisfied. Suppose $\lambda = a/b$ and $\mu = c/d$ with $a, b, c, d \in \mathbb{Z}^*$, $\gcd(a, b) = 1$, and $\gcd(c, d) = 1$.

Suppose we want to evaluate the Tutte polynomial for $H = (V_H, E_H)$ at x, y . Let $n := |V_H|$ and $m := |E_H|$. By (12), to evaluate $T(H; x, y)$, we can instead evaluate $Z(H; 1/\lambda - 1, \mu^2)$ (note that $\lambda \neq 1$ implies $x \neq 1$ and $y \neq 1$ and hence (12) applies). Recall that

$$Z(H; 1/\lambda - 1, \mu^2) = \sum_{S \subseteq E_H} (1/\lambda - 1)^{\kappa(S)} \mu^{2|S|} = \frac{L}{a^n d^{2m}}, \tag{29}$$

where

$$L = \sum_{S \subseteq E_H} (b-a)^{\kappa(S)} a^{n-\kappa(S)} c^{2|S|} d^{2m-2|S|}.$$

Note that $L \in \mathbb{Z}$ and $|L| \leq 2^m |b-a|^n a^n c^{2m} d^{2m}$.

We now prove the case $\mu \neq -2$. For the case $\mu = -2$, the proof is similar (by using Υ_2 and Corollary 2).

We choose n^3 primes $p_1, \dots, p_{n^3}$ such that

- $a, b, c, d \not\equiv 0 \pmod{p_i}$, and $a+b \not\equiv 0 \pmod{p_i}$, for each $i \in [n^3]$;
- there is some integer k for which (27) is satisfied with $p = p_i$, for each $i \in [n^3]$;
- $p_i = O(n^4)$ for $i \in [n^3]$; and
- $\prod_{i=1}^{n^3} p_i > 2^{m+1} |b-a|^n a^n c^{2m} d^{2m}$.

By Corollary 1, these primes exist. We can find them in time polynomial in n by exhaustive search.

For each p_i , let $0 < k_i < p_i$ be an integer for which (25) satisfies (17) with $p = p_i$ (by Fermat's little Theorem, if k_i is a solution, then $k_i + t(p_i - 1)$ is a solution as well, for every $t \in \mathbb{Z}$). Again, we can find them in time polynomial in n by exhaustive search.

We use Υ_1 with $k = k_i$ as above. Let G_i be the stretch-sum of H and (Υ_1, u_0) as above. Note that G_i has size polynomial in n since $k_i = O(n^4)$. By Lemma 7 and (29), we have

$$W \equiv a^n d^{2m} \lambda^{-2n} X^{-n} R'_2(G_i; \lambda, \mu) \pmod{p_i},$$

where $X = \lambda Z'_p(\Upsilon_1; \lambda, \mu)$. We can make a query to the oracle to obtain the rational number $R'_2(G_i; \lambda, \mu)$ and thus can compute $L \pmod{p_i}$ in polynomial time for each $i \in [n^3]$. By the Chinese remainder theorem, we can compute L in time polynomial in n (see, e.g., [3], p.106). $\square$

3.2 Reducing #PBIS to the R'_2 polynomial (proof of Theorem 5)

Now we show the connection between #PBIS and the R'_2 polynomial; the proof of Theorem 5 is similar to the proof of the high-temperature expansion of the Ising model (see, e.g., [26]).

Proof of Theorem 5.

$$\begin{aligned} \#PBIS(G; \eta) &= \sum_{\sigma: U \cup W \rightarrow \{0,1\}} (1+\eta)^{w(\sigma)} (1-\eta)^{|E|-w(\sigma)} \\ &= \sum_{\sigma: U \cup W \rightarrow \{0,1\}} \prod_{\{u,v\} \in E} (1 + \eta \chi(\sigma(u), \sigma(v))), \end{aligned} \tag{30}$$

where

$$\chi(\sigma(u), \sigma(v)) = \begin{cases} 1 & \text{if } \sigma(u) = \sigma(v) = 1 \\ -1 & \text{otherwise.} \end{cases} \tag{31}$$

Let

$$\Psi_{S, \sigma_1, \sigma_2} := \prod_{\{u,v\} \in S} \chi(\sigma_1(u), \sigma_2(v)) \quad \text{and} \quad \Psi_{S, \sigma_1} := \sum_{\sigma_2: W \rightarrow \{0,1\}} \Psi_{S, \sigma_1, \sigma_2}.$$

Expanding the product in (30) and changing the order of summation yields

$$\begin{aligned}
(30) &= \sum_{\sigma: U \cup W \rightarrow \{0,1\}} \sum_{S \subseteq E} \prod_{\{u,v\} \in S} \eta \chi(\sigma(u), \sigma(v)) \\
&= \sum_{S \subseteq E} \eta^{|S|} \sum_{\sigma: U \cup W \rightarrow \{0,1\}} \prod_{\{u,v\} \in S} \chi(\sigma(u), \sigma(v)) \\
&= \sum_{S \subseteq E} \eta^{|S|} \sum_{\sigma_1: U \rightarrow \{0,1\}} \Psi_{S, \sigma_1}. \tag{32}
\end{aligned}$$

Let $N_S(v)$ denote the set of neighbors of v in the subgraph $(U \cup W, S)$. Fix S and $\sigma_1: U \rightarrow \{0,1\}$. We say that a pair S, σ_1 is good if for every $v \in W$ the number of vertices $u \in N_S(v)$ such that $\sigma_1(u) = 1$ is even. A pair which is not good will be called bad.

Claim 2. $\Psi_{S, \sigma_1} = 2^{|W|}(-1)^{|S|}$ if the pair S, σ_1 is good; and $\Psi_{S, \sigma_1} = 0$ if the pair S, σ_1 is bad.

Proof of Claim 2. Suppose that σ_2 and σ'_2 differ only in the value assigned to $v \in W$. For every $u \in N_S(v)$ we have

$$\chi(\sigma_1(u), \sigma_2(v)) = \begin{cases} \chi(\sigma_1(u), \sigma'_2(v)) & \text{if } \sigma_1(u) = 0 \\ -\chi(\sigma_1(u), \sigma'_2(v)) & \text{if } \sigma_1(u) = 1. \end{cases}$$

Hence,

$$\Psi_{S, \sigma_1, \sigma_2} = \begin{cases} -\Psi_{S, \sigma_1, \sigma'_2} & \text{if } |\{u \in N_S(v) \mid \sigma_1(u) = 1\}| \text{ is odd,} \\ \Psi_{S, \sigma_1, \sigma'_2} & \text{otherwise.} \end{cases}$$

If the pair S, σ_1 is bad, then there is a vertex $v \in W$ such that $|\{u \in N_S(v) \mid \sigma_1(u) = 1\}|$ is odd. We can partition the $W \rightarrow \{0,1\}$ mappings into pairs σ_2, σ'_2 that differ only in the label of v . For each pair, we have $\Psi_{S, \sigma_1, \sigma_2} + \Psi_{S, \sigma_1, \sigma'_2} = 0$, and thus $\Psi_{S, \sigma_1} = 0$.

If the pair S, σ_1 is good, then each $\Psi_{S, \sigma_1, \sigma_2}$ contributes the same value $(-1)^{|S|}$. Since there are $2^{|W|}$ mappings from W to $\{0,1\}$ we have $\Psi_{S, \sigma_1} = 2^{|W|}(-1)^{|S|}$. $\square$

Claim 3. Fix $S \subseteq E$, the number of $\sigma_1: U \rightarrow \{0,1\}$ such that the pair S, σ_1 is good is $2^{|U| - \text{rk}_2(S)}$, where $\text{rk}_2(S)$ is the rank (over $\mathbb{F}_2$) of the bipartite adjacency matrix of $(U \cup W, S)$.

Proof of Claim 3. Let B be the bipartite adjacency matrix of $(U \cup W, S)$. Note that the pair S, σ_1 is good if and only if

$$\sigma_1^T B \equiv 0 \pmod{2}$$

(we view σ_1 as a vector with $(\sigma_1)_v = \sigma_1(v)$). The claim follows from the fact that the number of vectors $\alpha \in \{0,1\}^{|U|}$ such that $\alpha^T B \equiv 0 \pmod{2}$ is $2^{|U| - \text{rk}_2(B)}$. $\square$

By Claim 2 and Claim 3, (32) equals

$$\sum_{S \subseteq E} 2^{|U| + |W| - \text{rk}_2(S)} (-\eta)^{|S|} = 2^{|U| + |W|} R'_2(G; 1/2, -\eta).$$

$\square$

3.3 Reducing #BIS to #PBIS (proof of Lemma 4)

We use the following construction in the proof of Lemma 4. Given a graph $G = (V, E)$, a prime p and a positive integer k we construct a bipartite graph G' by replacing each vertex $v \in V$ by a cloud of kp new vertices and each edge $e \in E$ by a cloud of $p - 1$ new vertices. For every vertex $v \in V$ and every edge e adjacent to v we add a complete bipartite graph between the cloud of v and the cloud of e .

Lemma 9. *Let $p > 2$ be a prime, k be a positive integer, and η be a rational number such that $\eta \in \mathbb{Z}_p^*$ and*

$$\begin{aligned} \left(\frac{1+\eta}{1-\eta}\right)^{2k} + 1 &\equiv 0 \pmod{p}, \\ \eta - 1 &\not\equiv 0 \pmod{p}. \end{aligned} \quad (33)$$

Let $G = (V, E)$ be a graph and let $G' = (V', E')$ be the graph constructed above (using p, k). Then the number of independent sets of G is congruent to $\#PBIS(G'; \eta) \pmod{p}$.

Proof. We are going to evaluate

$$\#PBIS(G'; \eta) = \sum_{\sigma: V' \rightarrow \{0,1\}} \prod_{\{u,v\} \in E'} (1 + \eta \chi(\sigma(u), \sigma(v))) \quad (34)$$

modulo p (where χ is given by (31)).

Let s be a vertex of G . Consider a set of configurations (that is, σ 's in (34)) that differ only in the labels assigned to the vertices in the cloud of s (thus the configurations agree in the labels assigned to the vertices outside of the cloud of s). Note that by symmetry the value of $\prod_{\{u,v\} \in E'} (1 + \eta \chi(\sigma(u), \sigma(v)))$ depends only on h , the number of vertices in the cloud of s that are assigned 1. There are $\binom{kp}{h}$ such configurations and hence the total contribution of configurations with $h \in \{1, \dots, kp-1\}$ to (34) is zero modulo p . Thus, we only need to consider the σ 's such that for every $s \in V$ the vertices in the cloud of s are assigned the same label.

For $\varsigma: V \rightarrow \{0,1\}$ let C_ς be the set of assignments $V' \rightarrow \{0,1\}$ that assign label $\varsigma(s)$ to every vertex in the cloud of s (for every $s \in V$). Now we evaluate

$$\sum_{\sigma \in C_\varsigma} \prod_{\{u,v\} \in E'} (1 + \eta \chi(\sigma(u), \sigma(v))). \quad (35)$$

Note that each vertex in the edge clouds influences its own set of terms in the product in (35). Thus the sum in (35) turns into the following product

$$(35) = \prod_{\{s,t\} \in E} \Psi(\varsigma(s), \varsigma(t)), \quad (36)$$

where

$$\Psi(x, y) = [(1 + \eta \chi(x, 0))^{kp} (1 + \eta \chi(y, 0))^{kp} + (1 + \eta \chi(x, 1))^{kp} (1 + \eta \chi(y, 1))^{kp}]^{p-1}. \quad (37)$$

Now we evaluate $\Psi(x, y)$ for $x, y \in \{0, 1\}$

- Case 1: $x = y = 1$. We have

$$\Psi(1, 1) = [(1 - \eta)^{2kp} + (1 + \eta)^{2kp}]^{p-1},$$

and from the assumption (33) we obtain

$$\Psi(1, 1) = (1 - \eta)^{2kp(p-1)} \left[1 + \left(\frac{1 + \eta}{1 - \eta} \right)^{2kp} \right]^{p-1} \equiv 0 \pmod{p}. \quad (38)$$

- Case 2: $x = y = 0$. We have

$$\Psi(0, 0) = [2(1 - \eta)^{2kp}]^{p-1} \equiv 1 \pmod{p}, \quad (39)$$

where in the last congruence we used $\eta \not\equiv 1 \pmod{p}$ and Fermat's little Theorem.

- Case 3: $x = 1, y = 0$ (the case $x = 0, y = 1$ is the same). We have

$$\begin{aligned} \Psi(1, 0) &= [(1 - \eta)^{2kp} + (1 + \eta)^{kp}(1 - \eta)^{kp}]^{p-1} \\ &= (1 - \eta)^{2kp(p-1)} \left[1 + \left(\frac{1 + \eta}{1 - \eta} \right)^{kp} \right]^{p-1} \\ &\equiv 1 \pmod{p}, \end{aligned} \quad (40)$$

where the last congruence follows from Fermat's little Theorem and the fact that $((1 + \eta)/(1 - \eta))^k \not\equiv -1 \pmod{p}$ (since otherwise $((1 + \eta)/(1 - \eta))^{2k} \equiv 1 \pmod{p}$ which violates (33)).

From (38), (39), and (40) we obtain that (36) modulo p is 1 if ς corresponds to an independent set of G and is zero otherwise. Thus we have

$$\#\text{PBIS}(G'; \eta) \equiv \sum_{\varsigma \in V \rightarrow \{0,1\}} \prod_{\{s,t\} \in E} \Psi(\varsigma(s), \varsigma(t)) \equiv \#\text{BIS}(G) \pmod{p}.$$

□

Fix rational $\eta \notin \{\pm 1, 0\}$. We want to find sufficiently many primes p such that (33) can be satisfied for some integer k . The following results shows that the density (inside the set of all primes) of such primes is a positive constant.

Lemma 10 ([4]). *For $q \in \mathbb{Q}^*$ and $q \neq \pm 1$, the density (inside the set of all primes) of primes for which there exists $k \in \mathbb{Z}$ such that*

$$q^k \equiv -1 \pmod{p},$$

is a positive constant.

The mapping $\eta \mapsto (\frac{1+\eta}{1-\eta})^2$ maps $\eta \in \mathbb{Q}^* \setminus \{\pm 1\}$ to $\mathbb{Q}^* \setminus \{\pm 1\}$ and hence we have the following.

Corollary 3. *Fix rational $\eta \notin \{\pm 1, 0\}$. The density (inside the set of all primes) of the primes p such that (33) has an integer solution to k is a positive constant.*

Proof of Lemma 4. The proof is a routine application of the Chinese Remainder Theorem (similar to the proof of Lemma 3). Given an instance $G = (V, E)$ of $\#BIS$, let $n := |V|$ and L be the number of independent sets of G . Note that $L \leq 2^n$.

We choose n^2 primes $p_1, \dots, p_{n^2} > 2$ such that

- for each $i \in [n^2]$, (33) has an integer solution to k with $p = p_i$;
- $p_i = O(n^3)$ for $i \in [n^2]$, and
- $\prod_{i=1}^{n^2} p_i > 2^n$.

By Corollary 3, these primes exist when $\eta \notin \{\pm 1, 0\}$, and we can find them in time polynomial in n using exhaustive search.

For each p_i , let $0 < k_i < p_i$ be an integer solution of (33) with $p = p_i$ (if k_i is a solution of (33) then $k_i + t(p_i - 1)$ is a solution for every $t \in \mathbb{Z}$), again we can find the k_i using exhaustive search. We then construct a bipartite graph G'_i as above from G , p_i and k_i . Note that G'_i has size polynomial in n since $k_i < p_i = O(n^3)$. As (33) is satisfied, by Lemma 9, we have

$$L \equiv \#PBIS(G'_i; \eta) \pmod{p_i}.$$

We can make a query to the oracle to obtain the rational number $\#PBIS(G'_i; \eta)$ and thus can compute $L \pmod{p_i}$ in polynomial time for each $i \in [n^2]$. By the Chinese remainder theorem, we can compute L in time polynomial in n . $\square$

4 The linear-width of a graph

For the proof of Theorem 3 we will use the linear-width of a graph, a concept which was first defined by Thomas [37]. In this section we prove a bound on linear-width in terms of tree-width.

The *linear-width* of a graph $G = (V, E)$ is the smallest integer ℓ such that the edges of G can be arranged in a linear order $e_1, \dots, e_m$ in such a way that, for every $i \in [m]$, there are at most ℓ vertices that have an adjacent edge in $\{e_1, \dots, e_{i-1}\}$ and an adjacent edge in $\{e_i, \dots, e_m\}$. It is known that computing the linear-width of a graph is NP-complete [36]. For paths and cycles the linear-width is easy to compute.

Example 1. *The linear-width of a path is 1. The linear-width of a cycle is 2.*

Let $e_1, \dots, e_m$ be a permutation of the edges of $G = (V, E)$. We say that a vertex $v \in V$ is *dangerous* w.r.t. $i \in [m]$, if there exist two edges e_j, e_k adjacent to v such that $j < i \leq k$. Let D_i be the set of vertices which are dangerous w.r.t. i . Note that the linear-width of G is the minimum value of $\max_i |D_i|$ optimized over all permutations of the edges.

Now we give an upper bound on the linear-width for trees.

Lemma 11. *Let $T = (V, E)$ be a tree on n vertices. The linear-width of T is at most $\lfloor \log_2 n \rfloor$.*

Proof. We describe an ordering σ witnessing that the linear-width is at most $\lfloor \log_2 n \rfloor$. Pick an arbitrary vertex $r \in V$ to be the root. For a vertex v , let T_v be the subtree rooted at v and let $|T_v|$ be the number of vertices in T_v . We do a depth-first search (DFS) on T where the children are explored in the increasing order of $|T_w|$ (that is, the smaller subtrees are explored first). Let σ be the order in which the edges are discovered by the DFS.

Next, we show that the linear-width of $\sigma = e_1, \dots, e_m$ is at most $\lfloor \log_2 n \rfloor$. Let $i \in [m]$. Let $v_1, \dots, v_j$ be the vertices on the stack right before the edge e_i was discovered (thus e_i is adjacent to v_j). Note that the vertices which are not on the stack are not dangerous w.r.t. i (they either are already fully explored or haven't been discovered yet). Note that if v_{k+1} is the rightmost child of v_k then all the edges adjacent to v_k are discovered and hence v_k is not dangerous w.r.t. i (for all $k \in [j-1]$). On the other hand if v_{k+1} is not the rightmost child of v_k then our exploration order implies

$$|T_{v_{k+1}}| < |T_{v_k}|/2. \quad (41)$$

Note the following two facts: v_j may be dangerous and $|T_{v_j}| \geq 2$. If there are ℓ dangerous vertices among $v_1, \dots, v_{j-1}$ then by (41) we have $|T_w| \geq 2 \cdot 2^\ell$ where w is the topmost dangerous vertex. Using $|T_w| \leq n$ we obtain the result. $\square$

For general graphs we will show a generalization of Lemma 11: a bound on the linear-width of G in terms of the tree-width of G . We now define tree-width (see, e.g., [27] for a nice treatment).

Given a graph $G = (V, E)$, a *tree decomposition* of G is a pair $(T, \{U_h\}_{h \in V_T})$ where $T = (V_T, E_T)$ is a tree and $U_h \subseteq V$ satisfy: (i) each edge of G is in at least one subgraph induced by U_h ; and (ii) for any three vertices t_1, t_2, t_3 of T such that t_2 is in the path between t_1 and t_3 in T we have $U_{t_1} \cap U_{t_3} \subseteq U_{t_2}$. The *width* of a decomposition is $\max_{h \in V_T} |U_h| - 1$. The *tree-width* of G (denoted $\text{tw}(G)$) is the minimum width optimized over all tree decompositions.

Lemma 12. *Let $G = (V, E)$ be a graph. Then*

$$\text{linear-width}(G) \leq (\text{tw}(G) + 1)(\lfloor \log_2 n \rfloor + 1).$$

Proof. Given a graph G with $|V| = n$, let $(T, \{U_h\}_{h \in V_T})$ (where $T = (V_T, E_T)$) be an optimal tree decomposition of G . Let $r := |V_T|$. We can assume that $r \leq n$ (see, e.g., [27] p. 579). Let σ' be the order in which the vertices of T are discovered by the DFS of Lemma 11. W.l.o.g. let $\sigma' = 1, \dots, r$.

We will need the following “separation” property of dangerous vertices: a path between an explored and an unexplored vertex must contain a dangerous vertex (this follows from the fact that an explored vertex with an unexplored neighbor is dangerous).

Now we construct the ordering $\sigma = e_1, \dots, e_m$ on the edges of G . For each $h \in V_T$, let G_h be the induced subgraph of G on U_h . An edge $e \in E$ may appear in several G_h ; we “assign” it to the first G_h (in the order σ'). Let σ be an ordering such that edges assigned to G_h come before the edges assigned to $G_{h'}$ for any $h < h'$. Note that we do not impose any particular order on the edges assigned to G_h (for any $h \in [r]$).

Let $i \in [m]$ and let h be such that e_i is assigned to G_h . Let $D_i \subseteq V$ be the set of dangerous vertices w.r.t. i . Let $D'_h \subseteq V_T$ be the set of dangerous vertices just after h was discovered. Let $v \in D_i \setminus U_h$. We will show that $v \in U_g$ for some $g \in D'_h$.

Let e_{i_1} and e_{i_2} be adjacent to v and such that $i_1 < i \leq i_2$. Note that e_{i_1} and e_{i_2} cannot be assigned to G_h (since v is not in U_h). Thus e_{i_1} is assigned to an explored vertex j_1 of T and e_{i_2} is assigned to an unexplored vertex j_2 of T . Note that v is in U_{j_1} and U_{j_2} and hence it is in U_k for all k that are on the path between j_1 and j_2 in T . By the “separation” property of dangerous vertices one of them (say g) must be dangerous, and by the second property of tree decomposition U_g contains v .

Hence,

$$|D_i| \leq |U_h| + \sum_{g \in D'_h} |U_g| \leq (\text{tw}(G) + 1)(\lfloor \log_2 n \rfloor + 1).$$

$\square$

5 Analysis of the single bond flip chain for trees

Given a tree $G = (V, E)$, let Ω be the set of $2^{|E|}$ subsets of E . By Lemma 2, for every $H \subseteq E$, we know that $\text{rk}_2(H)$ is the size of maximum matching of the subgraph (V, H) . Let $w(H)$ be the size of maximum matching in a graph (V, H) . Let P be the transition matrix of the single bond flip Markov chain $\mathcal{M}$ from definition 2. It's easy to see that $\mathcal{M}$ is ergodic with unique stationary distribution π such that

$$\pi(H) \propto \lambda^{w(H)} \mu^{|E|}.$$

The goal of this section is to prove Theorem 3.

5.1 The canonical paths

We will bound the mixing time of our chain $\mathcal{M}$ using the canonical paths method, introduced in [8, 34, 25]. Now we go over the basic definitions for Markov chains, see, e. g., [24] for a comprehensive background.

Definition 6. *The total variation distance of two probability distribution ν and ν' on Ω is*

$$\|\nu - \nu'\|_{TV} = \frac{1}{2} \sum_{H \in \Omega} |\nu(H) - \nu'(H)| = \max_{S \subseteq \Omega} |\nu(S) - \nu'(S)|.$$

Definition 7. *The mixing time from initial state H , $\tau_H(\varepsilon)$, is defined as*

$$\tau_H(\varepsilon) = \min\{t : \|P^t(H, \cdot) - \pi\|_{TV} \leq \varepsilon\},$$

and the mixing time $\tau(\varepsilon)$ of the chain is defined as

$$\tau(\varepsilon) = \max_{H \in \Omega} \{\tau_H(\varepsilon)\}.$$

Let $\sigma = e_1, \dots, e_m$ be an ordering of the edges of $G = (V, E)$ (we will usually use the orderings supplied by Lemma 11 or Lemma 12). Given any pair $I, F \in \Omega$, let $I \oplus F$ be the symmetric difference of I and F (that is, the set of edges which are in either I or F but not in both). We define a canonical path $\gamma_{I,F}$ between I and F as follows. Let $e_{i_1}, \dots, e_{i_k}$ be the edges from $I \oplus F$ ordered according to σ (that is, $i_1 < i_2 < \dots < i_k$). Let

$$\gamma_{I,F} = (H_0, H_1, \dots, H_k), \tag{42}$$

where $H_0 = I$, $H_k = F$ and $H_j = H_{j-1} \oplus \{e_{i_j}\}$.

Lemma 13. *Let $G = (V, E)$ be a graph. Let $\sigma = e_1, \dots, e_m$ be an ordering on E with linear-width ℓ . Let I, F be subsets of E and let H be on the canonical path (42) (that is, $H = H_j$ for some $j \in \{0, \dots, k\}$). Then*

$$|w(I) + w(F) - w(H) - w(C)| \leq \ell,$$

where $C = I \oplus F \oplus H$.

Proof. Let $Q = \{e_1, \dots, e_{i_j}\}$. Note that $H = (F \cap Q) \cup (I \cap Q^c)$, where Q^c is the complement of Q (that is, $E \setminus Q$). Similarly, $C = (I \cap Q) \cup (F \cap Q^c)$.

Let D be the set of dangerous vertices w.r.t. e_{ij+1} . Let M_I and M_F be maximum matchings of I and F , respectively. Let

$$M_H = (M_F \cap Q) \cup (M_I \cap Q^c) \quad \text{and} \quad M_C = (M_I \cap Q) \cup (M_F \cap Q^c).$$

Note that all vertices of M_H with degree ≥ 2 are in D (a vertex which is not D has all adjacent edges (in G) from Q or from Q^c and hence the adjacent edges (in M_H) agree with M_I or M_F). The same is true for M_C . Moreover if a vertex $v \in D$ has degree 2 in M_H then it has degree 0 in M_C . Thus by removing $\leq |D|$ edges from M_H and M_C we can turn both of them into matchings. Thus

$$w(H) + w(C) \geq w(I) + w(F) - |D| \geq w(I) + w(F) - \ell. \quad (43)$$

Note that a canonical path from $I' := H$ to $F' := C$ passes through $H' := I$ (with $C' := I' \oplus F' \oplus H' = F$). Thus

$$w(I) + w(F) = w(H') + w(C') \geq w(I') + w(F') - \ell = w(H) + w(C) - \ell. \quad (44)$$

Combining (43) and (44) we get the lemma. $\square$

5.2 The congestion of $\mathcal{M}$

Now we analyze the congestion of the collection $\Gamma = \{\gamma_{I,F} \mid I, F \in \Omega\}$ where $\gamma_{I,F}$ are canonical paths defined in (42). For each transition (H, H') such that $P(H, H') > 0$, let $cp(H, H')$ be the set of pairs (I, F) such that $(H, H') \in \gamma_{I,F}$. The congestion of Γ on (H, H') is (see, e. g., [24])

$$\varrho_{(H,H')} = \frac{1}{P(H, H')} \sum_{I,F:(H,H') \in \gamma_{I,F}} \frac{\pi(I)\pi(F)}{\pi(H)} |\gamma_{I,F}|, \quad (45)$$

where $|\gamma_{I,F}|$ is the length of $\gamma_{I,F}$. The congestion of Γ is defined as

$$\varrho := \max_{\substack{(H,H'):\\ P(H,H') > 0}} \varrho_{(H,H')}.$$

We will use the following connection between the congestion and the mixing time.

Theorem 6 ([8, 34]). $\tau_H(\varepsilon) \leq \varrho(\log(1/\pi(H)) + \log(1/\varepsilon))$ for each starting state $H \in \Omega$.

At the end of this section we prove the following bound on the congestion of Γ .

Lemma 14. *Let $G = (V, E)$ be a graph. Let $\sigma = e_1, \dots, e_m$ be an ordering on E with linear-width ℓ . For every (H, H') such that $P(H, H') > 0$, and for every $\lambda, \mu > 0$ we have*

$$\varrho_{(H,H')} \leq 2|E|^2 \bar{\lambda}^\ell,$$

where $\bar{\lambda} = \max\{\lambda, 1/\lambda\}$.

We can now prove Theorem 3.

Proof of Theorem 3. Since $G = (V, E)$ is a tree, by Lemma 11, we have $\ell \leq \lfloor \log_2 n \rfloor$, by Lemma 14, we have

$$\varrho \leq 2|E|^2 \bar{\lambda}^\ell \leq 2|E|^2 n^{\lfloor \log_2 n \rfloor} \leq 2n^{2+\lfloor \log_2 n \rfloor}.$$

Theorem 3 now follows from Theorem 6. $\square$

Now we bound the congestion of our canonical paths.

Proof of Lemma 14. We will bound $\varrho_{(H,H')}$ for every (H, H') such that $P(H, H') > 0$. Let $\hat{H} = H$ if $\pi(H) \leq \pi(H')$ and $\hat{H} = H'$ otherwise. Note that

$$\frac{\pi(\hat{H})}{2|E|} = \pi(H)P(H, H') = \pi(H')P(H', H), \quad (46)$$

since $\mathcal{M}$ is reversible. We define a mapping $f : cp(H, H') \rightarrow \Omega$ such that $f(I, F) = I \oplus F \oplus \hat{H}$ for every pair $(I, F) \in cp(H, H')$.

First, note that f is an injection. Given $J \in \Omega$ we can determine the unique I, F such that $f(I, F) = J$, by first computing $J \oplus \hat{H}$, and then using the ordering σ on the edges of G to recover I and F .

Note that

$$|I| + |F| = |\hat{H}| + |f(I, F)|, \quad (47)$$

and

$$|w(I) + w(F) - w(\hat{H}) - w(f(I, F))| \leq \ell, \quad (48)$$

where (48) follows from Lemma 13.

Let $L = \sum_J \lambda^{w(J)} \mu^{|J|}$. We have the following upper bound on $\varrho_{(H,H')}$. By (45) and (46), we have

$$\begin{aligned} \varrho_{(H,H')} &= 2|E| \sum_{(I,F) \in cp(H,H')} \frac{\pi(I)\pi(F)}{\pi(\hat{H})} |\gamma_{I,F}| \\ &= 2|E|^2 \sum_{(I,F) \in cp(H,H')} \frac{\lambda^{w(I)+w(F)-w(\hat{H})} \mu^{|I|+|F|-|\hat{H}|}}{L} \\ &\leq 2|E|^2 \bar{\lambda}^\ell \sum_{(I,F) \in cp(H,H')} \frac{\lambda^{w(f(I,F))} \mu^{|f(I,F)|}}{L} \\ &\leq 2|E|^2 \bar{\lambda}^\ell, \end{aligned} \quad (49)$$

$$\leq 2|E|^2 \bar{\lambda}^\ell, \quad (50)$$

where (49) follows from (47) and (48), and (50) follows from the fact that f is an injection from $cp(H, H')$ to Ω . $\square$

6 The single bond flip chain for the random cluster model

The following sampling problem corresponds to the random cluster model defined in (1).

RANDOM CLUSTER MODEL with $q, \mu \geq 0$ ($\text{RC}(q, \mu)$)

Instance: a graph $G = (V, E)$.

Output: $S \subseteq E$ with probability of $S \propto q^{\kappa(S)} \mu^{|S|}$.

The random cluster model was introduced by Fortuin and Kasteleyn [15] as a model in statistical physics, and was intensively studied since then (see, e.g., the monograph by Grimmett [21]). By the results of [23, 42, 41], the problem of exactly computing the partition function of $\text{RC}(q, \mu)$ is

#P-hard even in bipartite planar graphs for all rational numbers $q, \mu > 0$ except when (i) $q = 1$; and (ii) $q = 2$.

In the context of approximate counting, little is known about $\#RC(q, \mu)$. The case $q = 2$ and $\mu > 0$ corresponds to the ferromagnetic Ising model and has an FPRAS [26]. For “dense graphs” an FPRAS exists for all $q, \mu \geq 0$, [2, 1].

Cooper and Frieze [7] and Cooper et al. [6] investigated the Swendsen-Wang algorithm for $RC(q, \mu)$. They proved, using coupling and conductance methods, that the Swendsen-Wang algorithm is rapidly mixing on some classes of graphs. Gore and Jerrum [20] pointed out that the single bond flip chain for random cluster model is torpid (mixes in exponential time) on the complete graph K_n for $q \geq 3$ and a value of μ depending on the size of the graph (that is, μ is not a fixed constant).

We will show that the single bond flip chain mixes for graphs with bounded tree-width. Our motivation is “showing that Markov chain mixes”, not approximation of $\#RC(q, \mu)$ (since the problem of computing the partition function of $RC(q, \mu)$ is polynomial-time solvable [31] for bounded tree-width graphs).

The single bond flip for the random cluster model is analogous to the chain from Definition 2. At time t , pick an edge $e \in E$ at random and let $S = X_t \oplus \{e\}$. Set $X_{t+1} = S$ with probability $(1/2) \min\{1, q^{\kappa(S) - \kappa(X_t)} \mu^{|S| - |X_t|}\}$ and $X_{t+1} = X_t$ with the remaining probability. The chain $\mathcal{M}$ defined above is ergodic with unique stationary distribution π such that $\pi(H) \propto q^{\kappa(H)} \mu^{|H|}$. Let P be the transition matrix of $\mathcal{M}$.

We will bound the mixing time of $\mathcal{M}$ for every fixed constant $q, \mu > 0$. Given an ordering σ on the edges of G , we define the same canonical paths for each pair of subgraphs I, F , and the same mapping f for every (H, H') such that $P(H, H') > 0$, as in Section 5. We present a result analogous to Lemma 13.

Lemma 15. *Let $G = (V, E)$ be a graph. Let $\sigma = e_1, \dots, e_m$ be an ordering on E with linear-width ℓ . Let I, F be subgraphs of E and let H be on the canonical path (42) (that is, $H = H_j$ for some $j \in \{0, \dots, k\}$). Then*

$$|\kappa(I) + \kappa(F) - \kappa(H) - \kappa(C)| \leq \ell,$$

where $C = I \oplus F \oplus H$.

Proof. As in the proof of Lemma 13 let $Q = \{e_1, \dots, e_{i_j}\}$. Let D be the set of dangerous vertices w.r.t. e_{i_j+1} . We can split $V \setminus D$ into $V_1 \cup V_2$, where V_1 contains vertices whose adjacent edges are from Q and V_2 contains vertices whose adjacent edges are from Q^c .

Let I_i be the subgraph of I induced by $V_i \cup D$ and F_i be the subgraph of F induced by $V_i \cup D$ (for $i = 1, 2$). Note that H can be viewed as a gluing of I_2 and F_1 on D ; we will denote this $H = I_2 \cup F_1$. Similarly $C = I_1 \cup F_2$.

First note that

$$\kappa(F_1) + \kappa(F_2) \leq \kappa(F) + |D|,$$

since F_1, F_2 can be obtained from F by splitting each vertex in D and the splitting of one vertex increases the number of components by at most one.

Let I' consist of the connected components of I that do not contain a vertex from D . Let I'_i be the subgraph of I induced by V_i (for $i = 1, 2$). Note that

$$\begin{aligned} \kappa(F_1 \cup I'_2) + \kappa(F_2 \cup I'_1) &= \kappa(F_1) + \kappa(I'_2) + \kappa(F_2) + \kappa(I'_1) = \\ \kappa(F_1) + \kappa(F_2) + \kappa(I') &\leq \kappa(F) + |D| + \kappa(I') \leq \kappa(F) + \kappa(I) + |D|, \end{aligned} \tag{51}$$

where the first and second inequalities follow from the fact that the following pairs of graphs are vertex disjoint: F_1 and I'_2 , F_2 and I'_1 , and I'_1 and I'_2 .

Note that $I_2 \oplus I'_2$ consists of connected components that contain a vertex from D . Thus $\kappa(F_1 \cup I_2) = \kappa(F_1 \cup I'_2)$ (since adding a component from $I_2 \oplus I'_2$ to $F_1 \cup I'_2$ only attaches that component to an existing component). Similarly $\kappa(F_2 \cup I_1) = \kappa(F_2 \cup I'_1)$. Combining with (51) we obtain

$$\kappa(H) + \kappa(C) \leq \kappa(I) + \kappa(F) + |D| \leq \kappa(I) + \kappa(F) + \ell. \quad (52)$$

Note that a canonical path from $I' := H$ to $F' := C$ passes through $H' := I$ (with $C' := I' \oplus F' \oplus H' = F$). Thus

$$\kappa(I) + \kappa(F) = \kappa(H') + \kappa(C') \leq \kappa(I') + \kappa(F') + \ell = \kappa(H) + \kappa(C) + \ell. \quad (53)$$

Combining (52) and (53) we get the lemma. $\square$

We have the following bound on the congestion.

Lemma 16. *Let $G = (V, E)$ be a graph. Let $\sigma = e_1, \dots, e_m$ be an ordering on E with linear-width ℓ . For every (H, H') such that $P(H, H') > 0$, and for every $q, \mu > 0$ we have*

$$\varrho_{(H, H')} \leq 2|E|^2 \bar{q}^\ell,$$

where $\bar{q} = \max\{q, 1/q\}$.

Proof. Same as the proof of Lemma 14. $\square$

By Theorem 6 and Lemma 16, we have the following result.

Theorem 7. *Given an graph $G = (V, E)$ with $m = |E|$, $n = |V|$ and linear-width ℓ , the mixing time $\tau(\varepsilon)$ of $\mathcal{M}$ is*

$$\tau(\varepsilon) = O\left(m^2 \bar{q}^\ell (m + n|\log q| + m|\log \mu| + \log(1/\varepsilon))\right),$$

where $\bar{q} = \max\{q, 1/q\}$.

By Lemma 12 we conclude the following.

Corollary 4. *Let k be an integer and let $q, \mu > 0$. Let $G = (V, E)$ be a graph with tree-width bounded by k , $n := |V|$, and $m := |E|$. The mixing time $\tau(\varepsilon)$ of the single bond flip chain $\mathcal{M}$ for $\text{RC}(q, \mu)$ is bounded as follows*

$$\tau(\varepsilon) = O\left(m^2 \bar{q}^{(k+1)(1+\log_2 n)} (m + n|\log q| + m|\log \mu| + \log(1/\varepsilon))\right),$$

where $\bar{q} = \max\{q, 1/q\}$.

We note that we do not need to find the linear-width (or tree-width) of a graph efficiently, the existence result is enough for our proof of the mixing time.

7 Conclusions

We conclude with an observation that a generalization of $\text{RWM}(\lambda, \mu)$ does not have an FPRAS (unless $\text{RP}=\text{NP}$) and a few questions.

Let A be an $m \times n$ matrix whose entries are zeros, ones, and indeterminates, where each indeterminate occurs once. A *completion* of A is a substitution of 0, 1 to all the indeterminates in A . We denote $\mathcal{C}_A$ to be the set of all completions of A . Let $\text{rk}_2(B)$ be the rank of B over $\mathbb{F}_2$. Can we sample B from $\mathcal{C}_A$ with the probability of B proportional to $\lambda^{\text{rk}_2(B)}$? Note that this problem is a generalization of the $\text{RWM}(\lambda, 1)$ problem. It turns out that finding the minimum rank completion of a matrix is NP-hard (Proposition 2.1, [32]) and hence a sampler is unlikely (unless $\text{NP}=\text{RP}$), since for $\lambda = 2^{-n^2}$ a random completion will be the minimum rank completion (with constant probability). The sampling problem could be easy for sufficiently large λ (the problem of finding maximum rank completion is in P, see, e. g., Section 4.1 of [22]).

Question 1. *What other interesting properties are encoded by the polynomial?*

Question 2. *Can one sample maximum rank completions of a matrix?*

References

- [1] Noga Alon, Alan Frieze, and Dominic Welsh. Polynomial time randomized approximation schemes for Tutte-Gröthendieck invariants: the dense case. *Random Structures Algorithms*, 6(4):459–478, 1995.
- [2] James D. Annan. A randomised approximation algorithm for counting the number of forests in dense graphs. *Combin. Probab. Comput.*, 3(3):273–283, 1994.
- [3] Eric Bach and Jeffrey Shallit. *Algorithmic number theory. Vol. 1.* Foundations of Computing Series. MIT Press, Cambridge, MA, 1996. Efficient algorithms.
- [4] Christian Ballot. Density of prime divisors of linear recurrences. *Mem. Amer. Math. Soc.*, 115(551):viii+102, 1995.
- [5] Rodney J. Baxter. *Exactly solved models in statistical mechanics*. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London, 1989. Reprint of the 1982 original.
- [6] Colin Cooper, Martin E. Dyer, Alan M. Frieze, and Rachel Rue. Mixing properties of the Swendsen-Wang process on the complete graph and narrow grids. *J. Math. Phys.*, 41(3):1499–1527, 2000. Probabilistic techniques in equilibrium and nonequilibrium statistical physics.
- [7] Colin Cooper and Alan M. Frieze. Mixing properties of the Swendsen-Wang process on classes of graphs. *Random Structures Algorithms*, 15(3-4):242–261, 1999. Statistical physics methods in discrete probability, combinatorics, and theoretical computer science (Princeton, NJ, 1997).
- [8] Persi Diaconis and Daniel Stroock. Geometric bounds for eigenvalues of Markov chains. *Ann. Appl. Probab.*, 1(1):36–61, 1991.
- [9] Martin Dyer, Alan Frieze, and Mark Jerrum. On counting independent sets in sparse graphs. *SIAM J. Comput.*, 31(5):1527–1541 (electronic), 2002.

- [10] Martin Dyer, Leslie Ann Goldberg, Catherine Greenhill, and Mark Jerrum. The relative complexity of approximate counting problems. *Algorithmica*, 38(3):471–500, 2004. Approximation algorithms.
- [11] Martin Dyer, Leslie Ann Goldberg, and Mark Jerrum. An approximation trichotomy for boolean $\#$ CSP. *Journal of Computer and System Sciences*, In Press, Corrected Proof, 2009.
- [12] Martin Dyer and Catherine Greenhill. On Markov chains for independent sets. *J. Algorithms*, 35(1):17–49, 2000.
- [13] Joanna Ellis-Monaghan and Criel Merino. Graph polynomials and their applications I: the Tutte polynomial. *arXiv*, 0803.3079, Jun 2008.
- [14] Joanna Ellis-Monaghan and Criel Merino. Graph polynomials and their applications II: interrelations and interpretations. *arXiv*, 0806.4699, Jun 2008.
- [15] Cees M. Fortuin and Piet W. Kasteleyn. On the random-cluster model. I. Introduction and relation to other models. *Physica*, 57:536–564, 1972.
- [16] Gudmund Skovbjerg Frandsen and Peter Frands Frandsen. Dynamic matrix rank. In *Automata, languages and programming. Part I*, volume 4051 of *Lecture Notes in Comput. Sci.*, pages 395–406. Springer, Berlin, 2006.
- [17] Leslie Ann Goldberg and Mark Jerrum. The complexity of ferromagnetic Ising with local fields. *Combin. Probab. Comput.*, 16(1):43–61, 2007.
- [18] Leslie Ann Goldberg and Mark Jerrum. Approximating the partition function of the ferromagnetic Potts model. *arXiv*, 1002.0986, Feb 2010.
- [19] Leslie Ann Goldberg, Mark Jerrum, and Mike Paterson. The computational complexity of two-state spin systems. *Random Structures Algorithms*, 23(2):133–154, 2003.
- [20] Vivek K. Gore and Mark R. Jerrum. The Swendsen-Wang process does not always mix rapidly. *J. Statist. Phys.*, 97(1-2):67–86, 1999.
- [21] Geoffrey Grimmett. *The random-cluster model*, volume 333 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2006.
- [22] Nicholas J. A. Harvey, David R. Karger, and Kazuo Murota. Deterministic network coding by matrix completion. In *Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 489–498 (electronic), New York, 2005. ACM.
- [23] François Jaeger, Dirk Vertigan, and Dominic J. A. Welsh. On the computational complexity of the Jones and Tutte polynomials. *Math. Proc. Cambridge Philos. Soc.*, 108(1):35–53, 1990.
- [24] Mark Jerrum. *Counting, sampling and integrating: algorithms and complexity*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2003.
- [25] Mark Jerrum and Alistair Sinclair. Approximating the permanent. *SIAM J. Comput.*, 18(6):1149–1178, 1989.

- [26] Mark Jerrum and Alistair Sinclair. Polynomial-time approximation algorithms for the Ising model. *SIAM J. Comput.*, 22(5):1087–1116, 1993.
- [27] Jon Kleinberg and Eva Tardos. *Algorithm Design*. Addison-Wesley Longman Publishing Co., Inc., Boston, MA, USA, 2005.
- [28] Michael Luby and Eric Vigoda. Fast convergence of the Glauber dynamics for sampling independent sets. *Random Structures Algorithms*, 15(3-4):229–241, 1999. Statistical physics methods in discrete probability, combinatorics, and theoretical computer science (Princeton, NJ, 1997).
- [29] Pieter Moree and Peter Stevenhagen. A two-variable Artin conjecture. *J. Number Theory*, 85(2):291–304, 2000.
- [30] Pieter Moree and Peter Stevenhagen. Prime divisors of the Lagarias sequence. *J. Théor. Nombres Bordeaux*, 13(1):241–251, 2001. 21st Journées Arithmétiques (Rome, 2001).
- [31] James G. Oxley and Dominic J. A. Welsh. Tutte polynomials computable in polynomial time. *Discrete Math.*, 109(1-3):185–192, 1992. Algebraic graph theory (Leibnitz, 1989).
- [32] René Peeters. Orthogonal representations over finite fields and the chromatic number of graphs. *Combinatorica*, 16(3):417–431, 1996.
- [33] John S. Provan and Michael O. Ball. The complexity of counting cuts and of computing the probability that a graph is connected. *SIAM J. Comput.*, 12(4):777–788, 1983.
- [34] Alistair Sinclair. Improved bounds for mixing rates of Markov chains and multicommodity flow. *Combin. Probab. Comput.*, 1(4):351–370, 1992.
- [35] Peter Stevenhagen. Prime densities for second order torsion sequences. 2000. Preprint.
- [36] Dimitrios M. Thilikos. Algorithms and obstructions for linear-width and related search parameters. *Discrete Appl. Math.*, 105(1-3):239–271, 2000.
- [37] Robin Thomas. Tree-decompositions of graphs (lecture notes), 1996.
- [38] William T. Tutte. A ring in graph theory. *Proc. Cambridge Philos. Soc.*, 43:26–40, 1947.
- [39] William T. Tutte. A contribution to the theory of chromatic polynomials. *Canadian J. Math.*, 6:80–91, 1954.
- [40] Salil P. Vadhan. The complexity of counting in sparse, regular, and planar graphs. *SIAM J. Comput.*, 31(2):398–427 (electronic), 2001.
- [41] Dirk Vertigan. The computational complexity of Tutte invariants for planar graphs. *SIAM J. Comput.*, 35(3):690–712 (electronic), 2005.
- [42] Dirk Vertigan and Dominic J. A. Welsh. The computational complexity of the Tutte plane: the bipartite case. *Combin. Probab. Comput.*, 1(2):181–187, 1992.
- [43] Dror Weitz. Counting independent sets up to the tree threshold. In *STOC’06: Proceedings of the 38th Annual ACM Symposium on Theory of Computing*, pages 140–149. ACM, New York, 2006.

- [44] Dominic J. A. Welsh. *Complexity: knots, colourings and counting*, volume 186 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1993.
- [45] Mingji Xia, Peng Zhang, and Wenbo Zhao. Computational complexity of counting problems on 3-regular planar graphs. *Theoret. Comput. Sci.*, 384(1):111–125, 2007.