IMPORTANT: Write each problem on a separate sheet of paper. Write your name on each sheet. (Why? To speed up the grading we will use parallelization—each problem will be graded by a different TA. At the beginning of the class (on the date when the homework is due) there will be a separate pile for each problem.)

Homework 4 due: Nov. 10 (Thursday); collected before class in B&L 109.
Problem sessions before Homework 4 is due:
Nov. 7 (Monday) 6:15pm - 7:15pm in Hylan 202
Nov. 8 (Tuesday) 6:15pm - 7:15pm in Goergen 109
Nov. 9 (Wednesday) 6:15pm - 7:15pm in Gavett 310
Nov. 9 (Wednesday) 7:40pm - 8:40pm in Hutchinson 473
Exam 3 is on: Nov. 17 (Thursday).
Problem sessions before Exam 3:
Nov. 14 (Monday) 6:15pm - 7:15pm in Goergen 109
Nov. 15 (Tuesday) 6:15pm - 7:15pm in Goergen 109
Nov. 16 (Wednesday) 6:15pm - 7:15pm in Gavett 310
Nov. 16 (Wednesday) 7:40pm - 8:40pm in Hutchinson 473

## 1 <u>Homework</u> - solve and turn in

4.1 (due Nov. 10) Krusty the Clown purchased a new super-efficient SUV with two engines. The first engine runs on a 2:1 mixture of gasoline and ethanol and gets 12 mpg. The second engine runs on a 1:2 mixture of gasoline and ethanol and gets 22 mpg. Krusty has 12 gallons of gasoline and 14 gallons of ethanol. How far can he get?

(a) Write a linear program for the problem. You are allowed to have only two variables in your linear program.

(b) Solve the linear program (you do not need to use the simplex method).

(c) Write down the dual of your linear program from part (a).

(d) Solve the dual program (you do not need to use the simplex method).

**4.2** (due Nov. 10) Give a mathematical expression (sums are allowed; linear programs are NOT allowed) for the value of the two person zero-sum game with the following  $n \times n$  payoff matrix

$$P = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & n \end{pmatrix}.$$

(More precisely, the matrix P is defined by  $P_{ii} = i$  and  $P_{ij} = 0$  for  $i \neq j$ ).

(Hint: Solve for small n first and then generalize. For example, for n = 2 solve the following linear program:

 $\min z$ 

$$\left(\begin{array}{cc}1&0\\0&2\end{array}\right)\left(\begin{array}{c}x_1\\x_2\end{array}\right) \leq \left(\begin{array}{c}z\\z\end{array}\right)$$

$$x_1 + x_2 = 1$$
  
 $x_1 \ge 0, x_2 \ge 0;$ 

we are using the convention that the entries in the matrix are the payoff for the row player.)

**4.3** (due Nov. 10) Let a be a constant and x be a variable. What is the value of the following linear program? Give a simple mathematical expression.

$$\min x \\ x \ge a \\ x \ge -a$$

Let a, b be constants and x, y be variables. What is the value of the following linear program? Give a simple mathematical expression.

$\min x + y$	
	$x \ge a$
	$x \ge -a$
	$y \ge b$
	$y \ge -b$

(If you don't know where to start—plug in concrete values for a and b (make sure to use negative numbers too), solve the resulting linear program (either by hand or using a solver) and formulate a conjecture based on the experiments.)

Let a, b be constants and x, y be variables. What is the value of the following linear program? Give a simple mathematical expression.

$$\min x - y$$

$$x \ge a$$

$$x \ge -a$$

$$y \ge b$$

$$y \ge -b$$

## 2 Bonus Homework - solve and turn in

**4.4** (due Nov. 10)

PART A:

We are given n pairs of real numbers  $(x_1, y_1), \ldots, (x_n, y_n)$ . We want to find real numbers a, b which minimize

$$\max_{i \in \{1,...,n\}} |ax_i + y_i - b|.$$

Write a linear program for this problem.

PART B:

 $(x_1, y_1), \ldots, (x_n, y_n)$ . We want to find real numbers a, b which minimize

$$\sum_{i \in \{1,\dots,n\}} |ax_i + y_i - b|.$$

Write a linear program for this problem.

## PART C:

We are given n pairs of real numbers  $(x_1, y_1), \ldots, (x_n, y_n)$ . We want to find real numbers a, b which minimize

$$\sum_{i \in \{1, \dots, n\}} (ax_i + y_i - b)^2.$$

Find a formula for a and b (you do NOT need linear programming for this one).