

RESOLUTION

Resolution + paramodulation provide a simple basis for mechanized deduction & Q.A.

- dominant technique since A. Robinson '65
 - basis of Prolog
1. Add denial of desired conclusion ψ to premises Δ , forming $\Delta \cup \{\neg\psi\}$
 2. Convert $\Delta \cup \{\neg\psi\}$ to clause form
 3. Repeatedly resolve (+factor +paramodulate) till \square is derived.

Simple example

1. All dogs bark

$$(\forall x (\text{Dog}(x) \rightarrow \text{Barks}(x)))$$

2. Snoopy is a dog

$$\text{Dog}(\text{Snoopy}) \quad \Delta$$

Prove: Something barks

$$(\exists x \text{ Barks}(x))$$

Denial: Nothing barks

$$\neg (\exists x \text{ Barks}(x)) \quad \neg \varphi$$

Conversion:

1. $\neg \text{Dog}(x) \vee \text{Barks}(x)$

convert \rightarrow to \vee , drop \forall

2. $\text{Dog}(\text{Snoopy})$


no change

3. $\neg \text{Barks}(x)$

change $\neg \exists$ to $\forall \neg$, drop \forall Refutation

$$\begin{array}{ll} r[2, 1a] & 4. \text{Barks}(\text{Snoopy}) \\ r[4, 3] & 5. \square \end{array}$$

$$\begin{array}{l} (x/\text{Snoopy}) \\ (x/\text{Snoopy}) \end{array}$$

As a graph:CONVERSION TO CLAUSE FORM

cf., CNF in satisfiability testing

1. Eliminate \Rightarrow and \Leftrightarrow

$$\varphi \Rightarrow \psi \longrightarrow \neg \varphi \vee \psi$$

$$\varphi \Leftrightarrow \psi \longrightarrow (\varphi \Rightarrow \psi) \wedge (\psi \Rightarrow \varphi)$$

2. Move \neg inward (5 rules)

$$\begin{array}{ccccc} \neg(\varphi \vee \psi) & , & \neg(\varphi \wedge \psi) & , & \neg\neg\varphi & , & \neg(\forall x \varphi) & , & \neg(\exists x \varphi) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \neg\varphi \wedge \neg\psi & & \neg\varphi \vee \neg\psi & & \varphi & & (\exists x \neg\varphi) & & (\forall x \neg\varphi) \end{array}$$

3. Rename duplicate bound variables

4. Skolemize (getting rid of \exists 's)5. Drop \forall 's

E.g., $(\exists x \text{ Robot}(x)) \longrightarrow$ Robot(c)
 give name to "the x that exists"

E.g., $(\forall x (\exists y \text{ Bigger-than}(y, x)))$

↓ make up a function that picks an appropriate y for each x

Bigger-than(f(x), x)

We can ... we drop V's
... sometimes finds all
... logical literals

Skol(φ) is satisfiable iff φ is satisfiable
 unsatisfiable iff φ is unsatisfiable

E.g., "For every pair of persons in the world, there is a 3rd person who speaks all of the languages of the first two"

$$\forall x \forall y \exists z. S(x, y, z)$$

<u>x</u>	<u>y</u>	<u>f(x, y)</u>
Smith	Lafrance	Trudeau
Smith	Glikov	Yevtushenko
Kohl	Yevtushenko	Yevtushenko
Trudeau	Yevtushenko	Nabokov
		etc.

$$S(x, y, f(x, y)) \quad \text{dropping } V: \text{ literal}$$

6. Move V inward (distribute V over \wedge)

$$(\varphi \wedge \psi) \vee \chi \longrightarrow (\varphi \vee \chi) \wedge (\psi \vee \chi)$$

$$\chi \vee (\varphi \wedge \psi) \longrightarrow (\chi \vee \varphi) \wedge (\chi \vee \psi)$$

7. Drop \wedge


E.g.

$$(\forall x [(\exists y [P(x, y) \vee Q(x, y)]) \Rightarrow (\exists y R(x, y))])$$

negate

yields a Skolem function of x , after renaming y to z

See if you can get 2 clauses,
 $\neg P(x, y) \vee R(x, f(x))$, $\neg Q(x, y) \vee R(x, f(x))$

Some terminologyNarrow resolution (without unification)

$$\frac{\psi \vee \psi, \neg \psi \vee x}{\psi \vee x} \quad \text{more gen: } \frac{\psi \vee \psi_1 \vee \dots \vee \psi_m, \neg \psi_1 \vee x_1 \vee \dots \vee x_n}{\psi_1 \vee \dots \vee \psi_m \vee x_1 \vee \dots \vee x_n}$$

$$\text{cf., a syllogism: } \frac{P \Rightarrow Q, Q \Rightarrow R}{P \Rightarrow R}$$

$$\begin{array}{ll} \text{Let } P \text{ be } \neg \psi & \neg \psi \Rightarrow \psi, \quad \psi \Rightarrow x \\ Q \text{ " } \psi & \neg \psi \Rightarrow x \\ R \text{ " } x & \text{or} \end{array}$$

$$\frac{\psi \vee \psi, \neg \psi \vee x}{\psi \vee x} \quad \text{as above}$$

soundness?

Narrow resolution, with unification:

$$\begin{array}{ccc} P(x, f(x), z) & P(A, f(A), z) & P(A, f(A), z) \\ \xrightarrow{x/A} P(A, y, w) & \xrightarrow{y/f(A)} P(A, y, w) & \xrightarrow{z/w} P(A, f(A), w) \end{array}$$

most general unifier (mgu) $(x/A)(y/f(A))(z/w)$
i.e., constraints $x = A$, $y = f(A)$, $z = w$

Example

$$1. \neg P(x, y, u) \vee \neg P(y, z, v) \vee \neg P(u, z, w) \vee \underline{P(x, v, w)}_{1d}$$

$$2. \neg P(i(x), E, w) \vee P(u, x, E) \quad \underline{2a}$$

Want to resolve 1d, 2a.

Rename x, u in (2) $\dots \rightarrow s, t$

$$1. \neg P(x, y, u) \vee \neg P(y, z, v) \vee \neg P(u, z, w) \vee \underline{P(x, v, w)}$$

$$2. \neg P(i(s), E, t) \vee P(t, s, E)$$

mgu: $(x/i(s))(v/E)(t/w)$
or (w/t)

Unify (not normally shown separately) :

- 1'. $\neg P(i(s), y, u) \vee \neg P(y, z, E) \vee \neg P(u, z, w) \vee \underline{P(i(s), E, w)}$
- 2'. $\neg P(i(s), E, w) \vee P(w, s, E)$

$r[1a, 2a]$:

3. $\neg P(i(s), y, u) \vee \neg P(y, z, E) \vee \neg P(u, z, w) \vee P(w, s, E)$

As part of a refutation graph :

1. $\neg P(x, y, w) \vee \neg P(y, z, v) \vee \neg P(u, z, w) \vee \underline{P(x, v, w)}$
2. $\neg P(i(s), E, t) \vee P(t, s, E)$
3. $\neg P(i(s), y, u) \vee \neg P(y, z, E) \vee \neg P(u, z, w) \vee P(w, s, E)$

Factoring

E.g., $P(x, A) \vee Q(x, y) \vee \underline{P(F(y), z)}$

$(x/F(y))(z/A)$

factor $P(F(y), A) \vee Q(F(y), y)$

E.g., 1. $\neg P(x, y, u) \vee \neg P(y, z, v) \vee \neg P(u, z, w) \vee P(x, v, w)$

$(u/x)(z/y)(w/u)$ N.B.

factor $f[1a, c]$: $\neg P(x, y, x) \vee \neg P(y, x, v) \vee P(x, v, x)$

Why we need factoring :

1. $P(x) \vee P(y)$
2. $\neg P(u) \vee \neg P(w)$

} clearly unsatisfiable!

3. $P(y) \vee \neg P(w)$ $r[1a, 2a]$

4. $\neg P(w) \vee \neg P(w')$ $r[2a, 3a]$, after renaming
etc.

can't derive \square !