

## A Less trivial resolution proof

1. No one likes anyone of whom he is afraid  
 $\forall x, y. A(x, y) \Rightarrow \neg L(x, y)$  (domain: people)

2. No one likes anyone who does not like him  
 $\forall x, y. \neg L(y, x) \Rightarrow \neg L(x, y)$

3. All paranoids are afraid of everyone except their friends  
 $H(x, y)$ :  $x$  has friend  $y$   
 $\forall x, y. (P(x) \wedge \neg H(x, y)) \Rightarrow A(x, y)$

4. There is a paranoid, and all of his friends are also paranoid  
 $\exists x. P(x) \wedge \forall y. H(x, y) \Rightarrow P(y)$

Conclusion:

There is someone whom all non-paranoids dislike  
 $\exists x \forall y. \neg P(y) \Rightarrow \neg L(y, x)$

Denial

5.  $\neg \exists x \forall y. \neg P(y) \Rightarrow \neg L(y, x)$

## Intuitive argument:

- From 3, all paranoids are afraid of everyone except their friends, so by 1 they don't like anyone except their friends, so by 2, everyone but a paranoid's friends dislike him
- But by 4 there is a paranoid, & so by the previous argument all but his friends dislike him; and since all his friends are paranoids, all non-paranoids dislike him

So, use 3, 1, 2, 4, 5

## Clause form

$$1. \neg A(x, y) \vee \neg L(x, y)$$

$$2. L(y, x) \vee \neg L(x, y)$$

$$3. \neg P(x) \vee H(x, y) \vee A(x, y)$$


$$4'. P(c)$$

$$4''. \neg H(c, y) \vee P(y)$$

$$5. \forall x \exists y. \neg (P(y) \vee \neg L(y, x))$$

$$\rightsquigarrow 5'. \neg P(f(x))$$

$$5''. L(f(x), x)$$



Corads: Study the mathematical example (group theory) in the notes.

Finding the m.g.u. of ( $\tau_1, \dots, \tau_n$ ), ( $\tau'_1, \dots, \tau'_n$ )

1. Standardize apart
2.  $mgu := ()$
3. Scan the lists in parallel, left to right
  - a. If end of lists is reached, return  $mgu$  (i.e., successful exit, even if  $mgu = ()$ )
  - b. If current symbols identical, continue at (a) with the next pair (i.e., advance pointers)
  - c. If neither symbol is a variable, return FAIL
  - d. One symbol is a variable, say  $v$ , and the other is the beginning of some term  $\tau$  (a variable, constant, or functional expression).  
 If  $v$  occurs in  $\tau$ , return FAIL "occurs-check"  
 else (i) push  $(v/\tau)$  onto  $mgu$   
 (ii) make this substitution throughout both lists, from the current pointers forward  
 (iii) move the pointers over the (now identical)  $\tau$  terms

e.g.,  $(x, F(x)), (y, y)$  can't be unified

Unification example

$$P(s, g(s), t, h(s, t), u, k(s, t, u))$$

$$\neg P(v, w, m(w), x, n(w, x), y)$$

Factoring example

$$P(y, x, A) \vee P(F(y), z, z) \vee P(F(A), B, x)$$

Formal statement of resolution (narrow)

$$\frac{\pi(\sigma_1, \dots, \sigma_k) \vee \alpha_1 \vee \dots \vee \alpha_m, \neg \pi(\tau_1, \dots, \tau_k) \vee \beta_1 \vee \dots \vee \beta_n}{\alpha'_1 \vee \dots \vee \alpha'_m \vee \beta'_1 \vee \dots \vee \beta'_n}$$

where  $(\sigma_1, \dots, \sigma_k), (\tau_1, \dots, \tau_k)$  are unifiable with mgu  $\delta$ ;  
 $\alpha'_i = \alpha_i \delta, \beta'_j = \beta_j \delta$ , etc.

$\pi(\dots), \neg \pi(\dots)$  need not be initial

Deductive Problem Solving & Q.A.

Cordell Green (IJCAI '69, Mach. Int. '69):

To solve a problem or answer a question,  
 1. prove that a solution/answer exists;  
 2. extract solution/answer from variable bindings of the proof.

Yes-no questions

Make "parallel" attempts to prove or disprove the proposition in question  
 $\leftrightarrow$  yes, no

$\varphi?$   $\longrightarrow$  refute  $\neg \varphi \longrightarrow$  yes  
 $\searrow$  refute  $\varphi \longrightarrow$  no

Reconsider Snoopy question: Does some x bark?

Wh- Questions ("fill in the blank(s)")

E.g.,

- 1.  $\forall x. At(Bob, x) \Rightarrow At(Carol, x)$  Carol is where-  
ever Bob is
- 2.  $At(Bob, Dance)$  Bob is at the dance

Q: Where is Carol?

Prove: There exists a <sup>thing</sup> place where Carol is

$\exists x At(Carol, x)$

Denial:  $\neg \exists x At(Carol, x)$

3. i.e.,  $\forall x. \neg At(Carol, x)$

Refutation & Ans - extraction

- 1.  $\neg At(Bob, x) \vee At(Carol, x)$  given
- 2.  $At(Bob, Dance)$  given
- 3.  $\neg At(Carol, x) \vee \underline{Ans(x)}$  denial
- 4.  $At(Carol, Dance)$   $r[1, 2]$
- 5.  $\square \vee \underline{Ans(Dance)}$   $r[3, 4]$

Good answers, bad answers

- E.g., 1.  $\forall x \exists y. Works-for(x, y)$  Everyone works for someone
- 2.  $Works-for(Bob, Carol)$  Bob works for Carol

Q:  $\exists x. Works-for(Bob, x)$  Who does Bob work for?

Denial:  $\forall x. \neg Works-for(Bob, x)$

First proof

- 1.  $Works-for(x, F(x))$  given
- 2.  $Works-for(Bob, Carol)$  given
- 3.  $\neg Works-for(Bob, x) \vee \underline{Ans(x)}$  denial
- 4.  $\square \vee \underline{Ans(F(Bob))}$   $r[1, 2]$   
Bob works for his employer 😊

Second proof

- 4'  $\square \vee \underline{Ans(Carol)}$   $r[2, 2]$   
better!

Generally: try to get Skolem-free answers by preferring proof steps that don't substitute Skolem constants/functions into the answer.  
But: there may only be Skolem answers —  
e.g., "Whom does every non-paranoid dislike?"