


Adaptive annealing: a near-optimal
connection between
sampling and counting

Daniel Štefankovič
(University of Rochester)


Santosh Vempala
Eric Vigoda
(Georgia Tech)



Adaptive annealing is near-optimal

If you want to count using MCMC then statistical physics is useful.

L. Langford
(Georgia Tech)



Outline

1. Counting problems

2. Basic tools: Chernoff, Chebyshev

**3. Dealing with large quantities
(the product method)**


4. Statistical physics

5. Cooling schedules (our work)


6. More...

Counting


independent sets




spanning trees




matchings



perfect matchings




k-colorings




Counting


independent sets




spanning trees




matchings



perfect matchings








k-colorings




Compute the number of

spanning trees



Compute the number of

spanning trees




Kirchhoff's Matrix Tree Theorem: $\det(\mathbf{D} - \mathbf{A})_{vv}$

$$\begin{matrix} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} & - & \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \\ \mathbf{D} & & \mathbf{A} \end{matrix}$$

$$\det \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$$

Compute the number of

spanning trees



G →

polynomial-time
algorithm




number of spanning trees of **G**


Counting




independent sets




spanning trees




matchings



perfect matchings




k-colorings




Compute the number of


independent sets
(hard-core gas model)



independent set of a graph = subset S of vertices, no two in S are neighbors



independent sets = 7



**independent set = subset S of vertices
no two in S are neighbors**

independent sets =

G_1 ○

G_2 ○—○

G_3 ○—○—○

G_{n-2} ○—○—○— ... —○


G_{n-1} ○—○—○— ... —○—○

G_n ○—○—○— ... —○—○—○

independent sets =

G_1				2
G_2				3
G_3				5
G_{n-2}		...		F_{n-1}
G_{n-1}		...		F_n
G_n		...		F_{n+1}


independent sets = 5598861



**independent set = subset S of vertices
no two in S are neighbors**

Compute the number of

independent sets



G →

polynomial-time
algorithm


?



number of independent sets of **G**

Compute the number of

independent sets



G



polynomial-time
algorithm





number of independent sets of **G**

(unlikely)



graph $G \mapsto$ # independent sets in G



#P-complete

#P-complete even for 3-regular graphs

(Dyer, Greenhill, 1997)

graph $G \mapsto$ # independent sets in G



approximation

randomization


graph $G \mapsto$ # independent sets in G



approximation

randomization

which is
more important?



graph $G \mapsto \#$ independent sets in G


My world-view:

(true) randomness is important conceptually but NOT computationally (i.e., I believe $P=BPP$).

approximation makes problems easier (i.e., I believe $\#P \neq BPP$)

important?

randomization



We would like to know Q

Goal: random variable Y such that

$$P((1-\varepsilon)Q \leq Y \leq (1+\varepsilon)Q) \geq 1-\delta$$

“ Y gives $(1\pm\varepsilon)$ -estimate”

We would like to know Q

Goal: random variable Y such that

$$P((1-\varepsilon)Q \leq Y \leq (1+\varepsilon)Q) \geq 1-\delta$$

(fully polynomial randomized approximation scheme):

FPRAS:

G, ε, δ



polynomial-time
algorithm



Y

Outline

1. Counting problems
2. Basic tools: Chernoff, Chebyshev
3. Dealing with large quantities
(the product method)
4. Statistical physics
5. Cooling schedules (our work)
6. More...

We would like to know Q

1. Get an unbiased estimator X , i. e.,

$$E[X] = Q$$

2. “Boost the quality” of X :

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

The Bienaymé-Chebyshev inequality



We would like to know Q

Goal: random variable Y such that
 $P((1-\varepsilon)Q \leq Y \leq (1+\varepsilon)Q) \geq 1-\delta$

“ Y gives $(1\pm\varepsilon)$ -estimate”

$P(Y \text{ gives } (1\pm\varepsilon)\text{-estimate})$

$$\geq 1 - \frac{V[Y]}{E[Y]^2} \frac{1}{\varepsilon^2}$$

The Bienaymé-Chebyshev inequality



We would like to know Q

Goal: random variable Y such that
 $P((1-\varepsilon)Q \leq Y \leq (1+\varepsilon)Q) \geq 1-\delta$

“Y gives $(1\pm\varepsilon)$ -estimate”

$P(Y \text{ gives } (1\pm\varepsilon)\text{-estimate})$

$$\geq 1 - \frac{V[Y]}{E[Y]^2} \frac{1}{\varepsilon^2}$$

squared coefficient
of variation SCV

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n} \implies \frac{V[Y]}{E[Y]^2} = \frac{1}{n} \frac{V[X]}{E[X]^2}$$

The Bienaymé-Chebyshev inequality

Let X_1, \dots, X_n, X be independent, identically distributed random variables,
 $Q = E[X]$. Let

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Then

$P(Y \text{ gives } (1 \pm \varepsilon)\text{-estimate of } Q)$

$$\geq 1 - \frac{V[X]}{n E[X]^2} \frac{1}{\varepsilon^2}$$

Chernoff's bound

Let X_1, \dots, X_n, X be independent, identically distributed random variables, $0 \leq X \leq 1$, $Q = E[X]$. Let

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Then

$P(Y \text{ gives } (1 \pm \varepsilon)\text{-estimate of } Q)$

$$\geq 1 - e^{-\varepsilon^2 \cdot n \cdot E[X] / 3}$$

The Bienaymé-Chebyshev inequality

Let X_1, \dots, X_n, X be independent, identically distributed random variables, $Q = E[X]$. Let

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Then

$P(Y \text{ gives } (1 \pm \varepsilon)\text{-estimate of } Q)$

$$\geq 1 - \frac{V[X]}{n E[X]^2} \frac{1}{\varepsilon^2}$$

Chernoff's bound

Let X_1, \dots, X_n, X be independent, identically distributed random variables, $0 \leq X \leq 1$, $Q = E[X]$. Let

$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Then

$P(Y \text{ gives } (1 \pm \varepsilon)\text{-estimate of } Q)$

$$\geq 1 - e^{-\varepsilon^2 \cdot n \cdot E[X] / 3}$$



$$n = \frac{V[X]}{E[X]^2} \frac{1}{\varepsilon^2} \frac{1}{\delta}$$

Number of samples to achieve precision ε with confidence δ .

$$n = \frac{1}{E[X]} \frac{3}{\varepsilon^2} \ln(1/\delta)$$

$$0 \leq X \leq 1$$





$$n = \frac{V[X]}{E[X]^2} \frac{1}{\varepsilon^2} \frac{1}{\delta}$$

BAD (with red arrow pointing to the $\frac{1}{\delta}$ term)

Number of samples to achieve precision ε with confidence δ .

$$n = \frac{1}{E[X]} \frac{3}{\varepsilon^2} \ln(1/\delta)$$

GOOD (with blue arrow pointing to $\ln(1/\delta)$)

BAD (with red arrow pointing to $\frac{1}{E[X]}$)

BAD (with red arrow pointing to $0 \leq X \leq 1$)




Median “boosting trick”

$$n = \frac{1}{E[X]} \frac{4}{\varepsilon^2}$$


$$Y = \frac{X_1 + X_2 + \dots + X_n}{n}$$

BY BIENAYME-CHEBYSHEV:

$$P(\bullet \in \text{red line}) \geq 3/4$$



Median trick – repeat $2T$ times



BY BIENAYME-CHEBYSHEV:

$$P(\text{blue circle} \in \text{red segment}) \geq 3/4$$

BY CHERNOFF:

$$P(\text{red segment} \text{ contains } > T \text{ out of } 2T \text{ blue circles}) \geq 1 - e^{-T/4}$$

$$P(\text{red segment} \text{ contains median}) \geq 1 - e^{-T/4}$$



$$n = \frac{V[X]}{E[X]^2} \frac{32}{\epsilon^2} \ln(1/\delta)$$

+ median trick

$$n = \frac{1}{E[X]} \frac{3}{\epsilon^2} \ln(1/\delta)$$

$$0 \leq X \leq 1$$

BAD



Creating “approximator” from X

ε = precision

δ = confidence

$$n = \Theta \left(\frac{V[X]}{E[X]^2} \frac{1}{\varepsilon^2} \ln(1/\delta) \right)$$



Outline

1. Counting problems
2. Basic tools: Chernoff, Chebyshev
3. Dealing with large quantities
(the product method)
4. Statistical physics
5. Cooling schedules (our work)
6. More...

(approx) counting \Leftrightarrow sampling

Valleau, Card'72 (physical chemistry), Babai'79 (for matchings and colorings), Jerrum, Valiant, V. Vazirani'86

the outcome of the JVV reduction:

random variables: $X_1 X_2 \dots X_t$
such that

1) $E[X_1 X_2 \dots X_t] = \text{“WANTED”}$

2) the X_i are easy to estimate

squared coefficient
of variation (SCV)

$$\frac{V[X_i]}{E[X_i]^2} = O(1)$$

(approx) counting \Leftrightarrow sampling

1) $E[X_1 X_2 \dots X_t] = \text{“WANTED”}$

2) the X_i are easy to estimate

$$\frac{V[X_i]}{E[X_i]^2} = O(1)$$

Theorem (Dyer-Frieze'91)

$O(t^2/\epsilon^2)$ samples ($O(t/\epsilon^2)$ from each X_i)


give

$1 \pm \epsilon$ estimator of “WANTED” with $\text{prob} \geq 3/4$



JVV for **independent sets**

GOAL: given a graph G , estimate the number of independent sets of G





$$\# \text{ independent sets} = \frac{1}{P(\text{graph})}$$

The same diamond-shaped graph as above, enclosed in large parentheses, representing the probability function P .

JVV for independent sets

$$P(A \cap B) = P(A)P(B|A)$$



$$X_i \in [0, 1] \text{ and } E[X_i] \geq 1/2 \quad \Rightarrow \quad \frac{V[X_i]}{E[X_i]^2} = O(1)$$

JVV for independent sets

$$P(A \cap B) = P(A)P(B|A)$$

(approx) counting \Leftrightarrow sampling

Valleau, Card'72 (physical chemistry), Babai'79 (for matchings and colorings), Jerrum, Valiant, V. Vazirani'86


the outcome of the JVV reduction:

random variables: $X_1 X_2 \dots X_t$
such that

1) $E[X_1 X_2 \dots X_t] = \text{"WANTED"}$

2) the X_i are easy to estimate

squared coefficient of variation (SCV) $\frac{V[X_i]}{E[X_i]^2} = O(1)$



X_1

X_2


X_3

X_4

$$X_i \in [0, 1] \text{ and } E[X_i] \geq \frac{1}{2} \quad \Rightarrow \quad \frac{V[X_i]}{E[X_i]^2} = O(1)$$

Self-reducibility for independent sets


$$P\left(\begin{array}{c} \textcircled{?} \\ \textcircled{} \textcircled{?} \\ \textcircled{?} \end{array}\right) = \frac{5}{7}$$



Self-reducibility for independent sets



$$P(\text{diamond with 3 nodes in question}) = \frac{5}{7}$$

$$P(\text{3x3 grid of diamonds}) = \frac{7}{5} \times P(\text{2x2 grid of diamonds})$$




Self-reducibility for independent sets


$$P(\text{graph with 3 nodes and 3 edges, 2 nodes marked with ?}) = \frac{5}{7}$$



$$= \frac{7}{5}$$




$$= \frac{7}{5}$$



Self-reducibility for independent sets



$$P(\text{graph with 3 nodes and 2 edges, 2 nodes marked with ?}) = \frac{3}{5}$$

$$\text{(graph with 5 nodes and 4 edges, 3 nodes marked with ?)} = \frac{5}{3} \text{(graph with 5 nodes and 4 edges, 2 nodes marked with ?)}$$




Self-reducibility for independent sets


$$P(\text{graph with 3 nodes, 2 edges, 2 nodes marked with ?}) = \frac{3}{5}$$




$$= \frac{5}{3}$$




$$= \frac{5}{3}$$




Self-reducibility for independent sets



$$= \frac{7}{5}$$




$$= \frac{7}{5} \frac{5}{3}$$




$$= \frac{7}{5} \frac{5}{3} \frac{3}{2} \left(\begin{array}{c} \bullet \\ \circ \end{array} \right) = 7$$

JVV: If we have a sampler oracle:




then FPRAS using $O(n^2)$ samples.

JVV: If we have a sampler oracle:



then FPRAS using $O(n^2)$ samples.

ŠVV: If we have a sampler oracle:



then FPRAS using $O^*(n)$ samples.

Application – independent sets

$O^*(|V|)$ samples suffice for counting

Cost per sample (Vigoda'01, Dyer-Greenhill'01)
time = $O^*(|V|)$ for graphs of degree ≤ 4 .

Total running time:

$O^*(|V|^2)$.

Other applications

matchings

$O^*(n^2m)$

(using Jerrum, Sinclair'89)

spin systems:

Ising model

$O^*(n^2)$ for $\beta < \beta_c$

(using Marinelli, Olivieri'95)

k-colorings

$O^*(n^2)$ for $k > 2\Delta$










(using Jerrum'95)

total running time








Outline

1. Counting problems
2. Basic tools: Chernoff, Chebyshev
3. Dealing with large quantities
(the product method)
4. **Statistical physics**
5. Cooling schedules (our work)
6. More...






easy = hot







hard = cold










Hamiltonian



2



1



0

Big set = Ω

Hamiltonian

$$H : \Omega \rightarrow \{0, \dots, n\}$$

Goal: estimate $|H^{-1}(0)|$

$$|H^{-1}(0)| = E[X_1] \dots E[X_t]$$

Distributions between **hot** and **cold**

β = inverse temperature

$\beta = 0 \Rightarrow$ **hot** \Rightarrow uniform on Ω

$\beta = \infty \Rightarrow$ **cold** \Rightarrow uniform on $H^{-1}(0)$

$$\mu_{\beta}(\mathbf{x}) \propto \exp(-H(\mathbf{x})\beta)$$

(Gibbs distributions)

Distributions between **hot** and **cold**

$$\mu_{\beta}(\mathbf{x}) \propto \exp(-H(\mathbf{x})\beta)$$

$$\mu_{\beta}(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$

Normalizing factor = partition function

$$Z(\beta) = \sum_{\mathbf{x} \in \Omega} \exp(-H(\mathbf{x})\beta)$$

Partition function

$$Z(\beta) = \sum_{\mathbf{x} \in \Omega} \exp(-H(\mathbf{x})\beta)$$

have: $Z(0) = |\Omega|$

want: $Z(\infty) = |H^{-1}(0)|$

Partition function - example

$$Z(\beta) = \sum_{\mathbf{x} \in \Omega} \exp(-H(\mathbf{x})\beta)$$


have: $Z(0) = |\Omega|$

want: $Z(\infty) = |H^{-1}(0)|$

$$Z(\beta) = \begin{aligned} & 1 e^{-4\beta} \\ & + 4 e^{-2\beta} \\ & + 4 e^{-1\beta} \\ & + 7 e^{-0\beta} \end{aligned}$$

$$Z(0) = 16$$


$$Z(\infty) = 7$$



Assumption:

we have a sampler oracle for μ_β

$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$



Assumption:

we have a sampler oracle for μ_β

$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$

$\mathbf{W} \sim \mu_\beta$

Assumption:

we have a sampler oracle for μ_β


$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$


$$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$$

Assumption:

we have a sampler oracle for μ_β


$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$

$$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$$

can obtain the following ratio:

$$E[X] = \sum_{\mathbf{s} \in \Omega} \mu_\beta(\mathbf{s}) X(\mathbf{s}) = \frac{Z(\alpha)}{Z(\beta)}$$

Our goal restated

Partition function

$$Z(\beta) = \sum_{\mathbf{x} \in \Omega} \exp(-H(\mathbf{x})\beta)$$

Goal: estimate $Z(\infty) = |\mathcal{H}^{-1}(0)|$

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \frac{Z(\beta_2)}{Z(\beta_1)} \dots \frac{Z(\beta_t)}{Z(\beta_{t-1})} Z(0)$$

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

Our goal restated

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \frac{Z(\beta_2)}{Z(\beta_1)} \dots \frac{Z(\beta_t)}{Z(\beta_{t-1})} Z(0)$$

Cooling schedule:

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

How to choose the cooling schedule?

minimize length, while satisfying

$$\frac{V[X_i]}{E[X_i]^2} = O(1) \quad E[X_i] = \frac{Z(\beta_i)}{Z(\beta_{i-1})}$$

Our goal restated

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \cdot \frac{Z(\beta_2)}{Z(\beta_1)} \cdot \frac{Z(\beta_3)}{Z(\beta_2)} \cdots$$

Cooling schedule:

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots$$

How to choose the

minimize length

Assumption:

we have a sampler oracle for μ_β

$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$

$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$

can obtain the following ratio:

$$E[X] = \sum_{\mathbf{s} \in \Omega} \mu_\beta(\mathbf{s}) X(\mathbf{s}) = \frac{Z(\alpha)}{Z(\beta)}$$

$$\frac{V[X_i]}{E[X_i]^2} = O(1) \quad E[X_i] = \frac{Z(\beta_i)}{Z(\beta_{i-1})}$$

Outline

1. Counting problems
2. Basic tools: Chernoff, Chebyshev
3. Dealing with large quantities
(the product method)
4. Statistical physics
5. Cooling schedules (our work)
6. More...

Parameters: **A** and **n**

$$Z(\beta) = \sum_{\mathbf{x} \in \Omega} \exp(-H(\mathbf{x})\beta)$$

$$Z(0) = \mathbf{A}$$

$$H: \Omega \rightarrow \{0, \dots, \mathbf{n}\}$$

$$Z(\beta) = \sum_{\mathbf{k}=0}^{\mathbf{n}} a_{\mathbf{k}} e^{-\beta \mathbf{k}}$$


$$a_{\mathbf{k}} = |H^{-1}(\mathbf{k})|$$

Parameters

$$Z(0) = A \quad H:\Omega \rightarrow \{0, \dots, n\}$$



	A	n
independent sets	2^V	E
matchings	$\approx V!$	V
perfect matchings	V!	V
k-colorings	k^V	E

Parameters





matchings = # ways of marrying them so that no unhappy couple

Parameters





matchings = # ways of marrying them so that no unhappy couple

Parameters



matchings = # ways of marrying them so that no unhappy couple

P



marry ignoring “compatibility”
hamiltonian = number of unhappy couples

Parameters

$$Z(0) = A \quad H:\Omega \rightarrow \{0, \dots, n\}$$

	A	n
independent sets	2^V	E
matchings	$\approx V!$	V
perfect matchings	V!	V
k-colorings	k^V	E

Previous cooling schedules

$$Z(0) = A \quad H:\Omega \rightarrow \{0, \dots, n\}$$

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

Cooling schedules of length

$$O(n \ln A)$$

$$O((\ln n) (\ln A))$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

Previous cooling s

$$Z(0) = A$$

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

Cooling schedules of length

$$O(n \ln A)$$

$$O((\ln n) (\ln A))$$

Our goal restated

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \frac{Z(\beta_2)}{Z(\beta_1)} \dots \frac{Z(\beta_t)}{Z(\beta_{t-1})} Z(0)$$

Cooling schedule:

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

How to choose the cooling schedule?

minimize length, while satisfying

$$\frac{V[X_i]}{E[X_i]^2} = O(1) \quad E[X_i] = \frac{Z(\beta_i)}{Z(\beta_{i-1})}$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

$$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$$

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

$$1/e \leq X \leq 1$$

$$\frac{V[X]}{E[X]^2} \leq \frac{1}{E[X]} \leq e$$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

$$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$$

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

$$Z(\infty) = a_0 \geq 1$$

$$Z(\ln A) \leq a_0 + 1$$

$$E[X] \geq 1/2$$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

$$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$$

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

$$E[X] \geq 1/2e$$

Previous cooling schedules

$1/n, 2/n, 3/n, \dots, (\ln A)/n, \dots, \ln A$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

Cooling schedules of length

$$O(n \ln A)$$

$$O((\ln n) (\ln A))$$

Our goal restated

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \frac{Z(\beta_2)}{Z(\beta_1)} \dots \frac{Z(\beta_t)}{Z(\beta_{t-1})} Z(0)$$

Cooling schedule:

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

How to choose the cooling schedule?

minimize length, while satisfying

$$\frac{V[X_i]}{E[X_i]^2} = O(1) \quad E[X_i] = \frac{Z(\beta_i)}{Z(\beta_{i-1})}$$

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

(Bezáková, Štefankovič,
Vigoda, V. Vazirani'06)

No better fixed schedule possible

$$Z(0) = A \quad H: \Omega \rightarrow \{0, \dots, n\}$$

THEOREM:

A schedule that works for all

$$Z_a(\beta) = \frac{A}{1+a} (1 + a e^{-\beta n})$$

(with $a \in [0, A-1]$)

has LENGTH $\geq \Omega((\ln n)(\ln A))$

Parameters

$$Z(0) = A \quad H: \Omega \rightarrow \{0, \dots, n\}$$

Our main result:

can get adaptive schedule
of length $O^* ((\ln A)^{1/2})$

Previously:

non-adaptive schedules
of length $\Omega^*(\ln A)$

Related work

can get adaptive schedule
of length $O^* ((\ln A)^{1/2})$

Lovász-Vempala

Volume of convex bodies in $O^*(n^4)$
schedule of length $O(n^{1/2})$

(non-adaptive cooling schedule, using specific properties
of the “volume” partition functions)

Existential part

Lemma:

for every partition function there exists
a cooling schedule of length $O^*((\ln A)^{1/2})$

there exists
~~can get adaptive schedule~~
of length $O^*((\ln A)^{1/2})$

Cooling schedule (definition refresh)

$$Z(\infty) = \frac{Z(\beta_1)}{Z(\beta_0)} \frac{Z(\beta_2)}{Z(\beta_1)} \dots \frac{Z(\beta_t)}{Z(\beta_{t-1})} Z(0)$$

Cooling schedule:

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

How to choose the cooling schedule?

minimize length, while satisfying

$$\frac{V[X_i]}{E[X_i]^2} = O(1) \quad E[X_i] = \frac{Z(\beta_i)}{Z(\beta_{i-1})}$$

Express **SCV** using **partition function**

(going from β to α) $E[X] = \frac{Z(\alpha)}{Z(\beta)}$


$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$

$$\frac{E[X^2]}{E[X]^2} = \frac{Z(2\alpha - \beta) Z(\beta)}{Z(\alpha)^2} \leq C$$

//

$$\frac{V[X]}{E[X]^2} + 1$$

$$\frac{E[X^2]}{E[X]^2} = \frac{Z(2\alpha-\beta) Z(\beta)}{Z(\alpha)^2} \leq C$$



$$f(\gamma) = \ln Z(\gamma)$$

$$(f(2\alpha-\beta) + f(\beta))/2 \leq (\ln C)/2 + f(\alpha)$$

$$\leq C' = (\ln C)/2$$

Properties of **partition functions**

$$f(\gamma) = \ln Z(\gamma)$$

f is decreasing

f is convex

$$f'(0) \geq -n$$

$$f(0) \leq \ln A$$

Properties of partition functions

$$f(\gamma) = \ln Z(\gamma)$$

f is decreasing

f is convex

$$f'(0) \geq -n$$

$$f(0) \leq \ln A$$

$$f(\beta) = \ln \sum_{k=0}^n a_k e^{-\beta k}$$

$$f'(\beta) = \frac{-\sum_{k=0}^n a_k k e^{-\beta k}}{\sum_{k=0}^n a_k e^{-\beta k}}$$

$$(\ln f)' = \frac{f'}{f}$$

GOAL: proving Lemma:
for every partition function there exists
a cooling schedule of length $O^*((\ln A)^{1/2})$

$$f(\gamma) = \ln Z(\gamma)$$

f is decreasing

f is convex

$$f'(0) \geq -n$$

$$f(0) \leq \ln A$$





either f or f'
changes a lot

Let $K := \Delta f$

Then

$$\Delta(\ln |f'|) \geq \frac{1}{K}$$






Let $K := \Delta f$

Then

$$\Delta(\ln |f'|) \geq \frac{1}{K}$$


$c := (a+b)/2$, $\Delta := b-a$
 have $f(c) = (f(a)+f(b))/2 - 1$

f is convex



$$\begin{aligned} (f(a) - f(c)) / \Delta &\leq f'(a) \\ (f(c) - f(b)) / \Delta &\geq f'(b) \end{aligned}$$

$$\frac{f'(b)}{f'(a)} \leq 1 - 1/\Delta f \leq e^{-\Delta f}$$



Let $K := \Delta f$


Then

$$\Delta(\ln |f'|) \geq \frac{1}{K}$$

$c := (a+b)/2, \Delta := b-a$


have $f(c) = (f(a)+f(b))/2 - 1$

f is convex




$$\begin{aligned} (f(a) - f(c)) / \Delta &\leq f'(a) \\ (f(c) - f(b)) / \Delta &\geq f'(b) \end{aligned}$$

$f:[a,b] \rightarrow \mathbb{R}$, convex, decreasing
can be “approximated” using



Technicality: getting to $2\alpha - \beta$



β α $2\alpha - \beta$

Express **SCV** using **partition function**


(going from β to α) $E[X] = \frac{Z(\alpha)}{Z(\beta)}$

$W \sim \mu_\beta \rightarrow X = \exp(H(W)(\beta - \alpha))$


$$\frac{E[X^2]}{E[X]^2} = \frac{Z(2\alpha - \beta) Z(\beta)}{Z(\alpha)^2} \leq C$$

\parallel
 $\frac{V[X]}{E[X]^2} + 1$


Technicality: getting to $2\alpha-\beta$



Technicality: getting to $2\alpha - \beta$



Technicality: getting to $2\alpha-\beta$



In In A
extra
steps

Existential \rightarrow Algorithmic

there exists
~~can get adaptive schedule~~
of length $O^* ((\ln A)^{1/2})$



can get adaptive schedule
of length $O^* ((\ln A)^{1/2})$

Algorithmic construction

Our main result:

using a sampler oracle for μ_β

$$\mu_\beta(\mathbf{x}) = \frac{\exp(-H(\mathbf{x})\beta)}{Z(\beta)}$$

we can construct a cooling schedule of length

$$\leq 38 (\ln A)^{1/2} (\ln \ln A) (\ln n)$$

Total number of oracle calls

$$\leq 10^7 (\ln A) (\ln \ln A + \ln n)^7 \ln(1/\delta)$$

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2$$

$$E[X] = \frac{Z(\alpha)}{Z(\beta)}$$

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2$$

$$E[X] = \frac{Z(\alpha)}{Z(\beta)}$$

X is “easy to estimate”

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2$$


$$E[X] = \frac{Z(\alpha)}{Z(\beta)}$$

we make progress (where $B_1 > 1$)

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2$$

$$E[X] = \frac{Z(\alpha)}{Z(\beta)}$$



need to construct a “feeler” for this

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2 \quad E[X] = \frac{Z(\alpha)}{Z(\beta)}$$
$$\uparrow \quad = \quad \frac{Z(\beta)}{Z(\alpha)} \frac{Z(2\beta - \alpha)}{Z(\alpha)}$$

need to construct a “feeler” for this

Algorithmic construction

current inverse temperature β

ideally move to α such that

$$B_1 \leq \frac{E[X^2]}{E[X]^2} \leq B_2$$

$$\uparrow = \frac{Z(\beta)}{Z(\alpha)} \frac{Z(2\beta - \alpha)}{Z(\alpha)}$$

bad "feeler"

$$E[X] = \frac{Z(\alpha)}{Z(\beta)}$$

need to construct a "feeler" for this

estimator for $\frac{Z(\beta)}{Z(\alpha)}$

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

For $W \sim \mu_\beta$ we have $P(H(W)=k) = \frac{a_k e^{-\beta k}}{Z(\beta)}$

estimator for $\frac{Z(\beta)}{Z(\alpha)}$

If $H(X)=k$ likely at both $\alpha, \beta \Rightarrow$ estimator

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

For $W \sim \mu_\beta$ we have $P(H(W)=k) = \frac{a_k e^{-\beta k}}{Z(\beta)}$

For $U \sim \mu_\alpha$ we have $P(H(U)=k) = \frac{a_k e^{-\alpha k}}{Z(\alpha)}$

estimator for $\frac{Z(\beta)}{Z(\alpha)}$

If $H(X)=k$ likely at both $\alpha, \beta \Rightarrow$ estimator

$$Z(\beta) = \sum_{k=0}^n$$



$$n = \frac{V[X]}{E[X]^2} \frac{1}{\epsilon^2} \frac{1}{\delta}$$

Number of samples to achieve precision ϵ with confidence δ .

For $W \sim \mu_\beta$

$$n = \frac{1}{E[X]} \frac{3}{\epsilon^2} \ln(1/\delta)$$

For $U \sim \mu_\alpha$

$$0 \leq X \leq 1$$



estimator for $\frac{Z(\beta)}{Z(\alpha)}$

For $W \sim \mu_\beta$ we have $P(H(W)=k) = \frac{a_k e^{-\beta k}}{Z(\beta)}$

For $U \sim \mu_\alpha$ we have $P(H(U)=k) = \frac{a_k e^{-\alpha k}}{Z(\alpha)}$

$$\frac{P(H(U)=k)}{P(H(W)=k)} e^{k(\alpha-\beta)} = \frac{Z(\beta)}{Z(\alpha)}$$

estimator for $\frac{Z(\beta)}{Z(\alpha)}$

For $W \sim \mu_\beta$ we have $P(H(W)=k) = \frac{a_k e^{-\beta k}}{Z(\beta)}$

For $U \sim \mu_\alpha$ we have $P(H(U)=k) = \frac{a_k e^{-\alpha k}}{Z(\alpha)}$

$$\frac{P(H(U)=k)}{P(H(W)=k)} e^{k(\alpha-\beta)} = \frac{Z(\beta)}{Z(\alpha)}$$

PROBLEM: $P(H(W)=k)$ can be too small

Rough estimator for $\frac{Z(\beta)}{Z(\alpha)}$

$$Z(\beta) = \sum_{k=0}^n a_k e^{-\beta k}$$

interval instead
of single value

For $W \sim \mu_\beta$ we have

$$P(H(W) \in [c, d]) = \frac{\sum_{k=c}^d a_k e^{-\beta k}}{Z(\beta)}$$

For $U \sim \mu_\alpha$ we have

$$P(H(W) \in [c, d]) = \frac{\sum_{k=c}^d a_k e^{-\alpha k}}{Z(\alpha)}$$

Rough estimator for $\frac{Z(\beta)}{Z(\alpha)}$

If $|\alpha - \beta| \cdot |d - c| \leq 1$ then

$$\frac{1}{e} \frac{Z(\beta)}{Z(\alpha)} \leq \frac{P(H(U) \in [c, d])}{P(H(W) \in [c, d])} e^{c(\alpha - \beta)} \leq e \frac{Z(\beta)}{Z(\alpha)}$$

We also need $P(H(U) \in [c, d])$
 $P(H(W) \in [c, d])$ to be large.

$$\frac{\sum_{k=c}^d a_k e^{-\alpha k}}{\sum_{k=c}^d a_k e^{-\beta k}} e^{c(\alpha - \beta)} = \frac{\sum_{k=c}^d a_k e^{-\alpha (k-c)}}{\sum_{k=c}^d a_k e^{-\beta (k-c)}}$$

We will:

Split $\{0, 1, \dots, n\}$ into $h \leq 4(\ln n)\sqrt{\ln A}$
intervals

$[0], [1], [2], \dots, [c, c(1+1/\sqrt{\ln A})], \dots$

for any inverse temperature β there
exists a interval with $P(H(W) \in I) \geq 1/8h$

We say that I is **HEAVY** for β

We will:

Split $\{0, 1, \dots, n\}$ in intervals

$[0], [1], [2], \dots, [c]$

Rough estimator for $\frac{Z(\beta)}{Z(\alpha)}$

If $|\alpha - \beta| \cdot |d - c| \leq 1$ then

$$\frac{1}{e} \frac{Z(\beta)}{Z(\alpha)} \leq \frac{P(H(U) \in [c, d])}{P(H(W) \in [c, d])} e^{c(\alpha - \beta)} \leq e \frac{Z(\beta)}{Z(\alpha)}$$

We also need $P(H(U) \in [c, d])$
 $P(H(W) \in [c, d])$ to be large.

$$\frac{\sum_{k=c}^d a_k e^{-\alpha k}}{\sum_{k=c}^d a_k e^{-\beta k}} e^{c(\alpha - \beta)} = \frac{\sum_{k=c}^d a_k e^{-\alpha (k-c)}}{\sum_{k=c}^d a_k e^{-\beta (k-c)}}$$

for any inverse temperature β there exists a interval with $P(H(W) \in I) \geq 1/8h$

We say that I is **HEAVY** for β

Algorithm

repeat

find an interval I which is heavy for
the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$

ANALYSIS:

either

- * make progress, or
- * eliminate the interval I
- * or make a “long move”

Algorithm

repeat

I is
heavy


find an interval I which is heavy for
the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$



distribution of $h(X)$ where $X \sim \mu_\beta$



I = a heavy interval at β

Algorithm

repeat

I is heavy

I is NOT heavy


find an interval I which is heavy for the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$



distribution of $h(X)$ where $X \sim \mu_\gamma$



no longer heavy at γ !

I = a heavy interval at β

Algorithm

repeat

I is heavy


I is heavy

I is NOT heavy


find an interval I which is heavy for the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$



distribution of $h(X)$ where $X \sim \mu_{\gamma}$



heavy at γ'

I = a heavy interval at β



Algorithm

repeat

find an interval I which is heavy for the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$



$I=[a,b]$

use binary search to find β^*

$$\alpha = \min\{1/(b-a), \ln A\}$$


Algorithm

repeat

find an interval I which is heavy for the current inverse temperature β

see how far I is heavy (until some β^*)

use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$



I is heavy
 β^*

$$\beta^* + 1/(2n)$$

I is NOT heavy

$$I = [a, b]$$

use binary search to find β^*

$$\alpha = \min\{1/(b-a), \ln A\}$$

How do you know that you can use binary search?

How do you know that you can use binary search?

I is NOT
heavy

I is
heavy

I is
heavy

I is NOT
heavy

Lemma: the set of temperatures for which I is h-heavy is an interval.

I is h-heavy at $\beta \iff P(h(X) \in I) \geq 1/8h$ for $X \sim \mu_\beta$

$$\sum_{k \in I} a_k e^{-\beta k} \geq \frac{1}{8h} \sum_{k=0}^n a_k e^{-\beta k}$$

How do you know that you can use binary search?

$$\sum_{k \in I} a_k e^{-\beta k} \geq \frac{1}{8h} \sum_{k=0}^n a_k e^{-\beta k}$$

$$x = e^{-\beta}$$

Descarte's rule of signs:

$$c_0 x^0 + c_1 x^1 + c_2 x^2 + \dots + c_n x^n$$



sign change

number of positive roots \leq number of sign changes

How do you know that you can use binary search?

$$\begin{aligned} & -1+x+x^2+x^3+\dots+x^n \\ & 1+x+x^{20} \end{aligned}$$

Descarte's rule of signs:

$$c_0 x^0 + c_1 x^1 + c_2 x^2 + \dots + c_n x^n$$



sign change

number of positive roots \leq number of sign changes

How do you know that you can use binary search?

$$\sum_{k \in I} a_k e^{-\beta k} \geq \frac{1}{8h} \sum_{k=0}^n a_k e^{-\beta k}$$

$$x = e^{-\beta}$$

Descarte's rule of signs:

$$c_0 x^0 + c_1 x^1 + c_2 x^2 + \dots + c_n x^n$$




sign change

number of positive roots \leq number of sign changes

$I=[a,b]$

I is heavy



I is heavy
 β^*

$\beta^* + 1/(2n)$

I is NOT heavy

Algorithm

repeat

find an interval I which is heavy for the current inverse temperature β



see how far I is heavy (until some β^*)



use the interval I for the feeler $\frac{Z(\beta)}{Z(\alpha)}$ $\frac{Z(2\beta-\alpha)}{Z(\alpha)}$

Rough estimator for $\frac{Z(\beta)}{Z(\alpha)}$

If $|\alpha-\beta| \cdot |d-c| \leq 1$ then

$$\frac{1}{e} \frac{Z(\beta)}{Z(\alpha)} \leq \frac{P(H(U) \in [c,d])}{P(H(W) \in [c,d])} e^{c(\alpha-\beta)} \leq e \frac{Z(\beta)}{Z(\alpha)}$$

We also need $P(H(U) \in [c,d])$

$P(H(W) \in [c,d])$ to be large

$$\frac{\sum_{k=c}^d a_k e^{-\alpha k}}{\sum_{k=c}^d a_k e^{-\beta k}} e^{c(\alpha-\beta)} = \frac{\sum_{k=c}^d a_k e^{-\alpha(k-c)}}{\sum_{k=c}^d a_k e^{-\beta(k-c)}}$$


can roughly compute ratio of

$$Z(\alpha)/Z(\alpha')$$

for $\alpha, \alpha' \in [\beta, \beta^*]$

if $|\alpha-\alpha'| \cdot |b-a| \leq 1$

$I=[a,b]$



1. success

2. eliminate interval



3. long move



find largest α such that

$$\frac{Z(\beta)}{Z(\alpha)} \frac{Z(2\beta - \alpha)}{Z(\alpha)} \leq C$$

can roughly
compute ratio of
 $Z(\alpha)/Z(\alpha')$
for $\alpha, \alpha' \in [\beta, \beta^*]$
if $|\alpha - \alpha'| \cdot |b - a| \leq 1$

input : A black-box sampler for $X \sim \mu_\beta$ for any $\beta \geq 0$, starting inverse temperature β_0 .
output: A cooling schedule for Z .

Bad $\leftarrow \emptyset$

print β_0

if $\beta_0 < \ln A$ **then**

$I \leftarrow \text{FIND-HEAVY}(\beta_0, \text{Bad})$

$w \leftarrow$ the width of I

$L \leftarrow \min\{\beta_0 + 1/w, \ln A\}$; (where $1/0 = \infty$)

$\beta^* \leftarrow$ binary search on $\beta^* \in [\beta_0, L]$ with precision $1/(2n)$, using predicate

 IS-HEAVY(β^*, I)

$\beta \leftarrow$ binary search on $\beta \in [\beta_0, (\beta^* + \beta_0)/2]$ with precision $1/(4n)$,
 using predicate $\text{EST}(I, \beta_0, \beta) \cdot \text{EST}(I, 2\beta - \beta_0, \beta) \leq 2000$

if $\beta < (\beta^* + \beta_0)/2$ **then**

 PRINT-COOLING-SCHEDULE(β) (optimal move)

else

if $\beta = L$ **then**

 PRINT-COOLING-SCHEDULE(β) (long move)

else

$\gamma \leftarrow (\beta^* - \beta_0)/2$

 print $\beta_0 + \gamma, \beta_0 + (3/2)\gamma, \beta_0 + (7/4)\gamma, \dots, \beta_0 + (2 - 2^{-\lceil \ln \ln A \rceil})\gamma$

 Bad \leftarrow Bad $\cup I$

 PRINT-COOLING-SCHEDULE(β^*) (interval move)

end

end

else

 print ∞

end

if we have sampler oracles for μ_β
then we can get adaptive schedule
of length $t=O^* \left((\ln A)^{1/2} \right)$

independent sets $O^*(n^2)$
(using Vigoda'01, Dyer-Greenhill'01)

matchings $O^*(n^2m)$
(using Jerrum, Sinclair'89)

spin systems:

Ising model $O^*(n^2)$ for $\beta < \beta_c$
(using Marinelli, Olivieri'95)

k-colorings $O^*(n^2)$ for $k > 2\Delta$
(using Jerrum'95)

Outline

1. Counting problems
2. Basic tools: Chernoff, Chebyshev
3. Dealing with large quantities
(the product method)
4. Statistical physics
5. Cooling schedules (our work)
6. More...

Outline

6. More...

a) proof of Dyer-Frieze

b) independent sets revisited

c) warm starts

Appendix – proof of:

1) $E[X_1 X_2 \dots X_t] = \text{“WANTED”}$

2) the X_i are easy to estimate

$$\frac{V[X_i]}{E[X_i]^2} = O(1)$$



Theorem (Dyer-Frieze'91)

$O(t^2/\epsilon^2)$ samples ($O(t/\epsilon^2)$ from each X_i)

give

$1 \pm \epsilon$ estimator of “WANTED” with $\text{prob} \geq 3/4$

How precise do the X_i have to be?

First attempt – term by term

Main idea:

$$\left(1 \pm \frac{\varepsilon}{t}\right) \left(1 \pm \frac{\varepsilon}{t}\right) \left(1 \pm \frac{\varepsilon}{t}\right) \dots \left(1 \pm \frac{\varepsilon}{t}\right) \approx 1 \pm \varepsilon$$

$$n = \Theta \left(\frac{V[X]}{E[X]^2} \frac{1}{\varepsilon^2} \ln(1/\delta) \right)$$

each term $\Omega(t^2)$ samples $\Rightarrow \Omega(t^3)$ total

How precise do the X_i have to be?

Analyzing SCV is better

(Dyer-Frieze'1991)

$$X = X_1 X_2 \dots X_t$$

squared coefficient
of variation (SCV)

GOAL: $SCV(X) \leq \varepsilon^2/4$

$P(X \text{ gives } (1 \pm \varepsilon)\text{-estimate})$

$$\geq 1 - \frac{V[X]}{E[X]^2} \frac{1}{\varepsilon^2}$$

How precise do the X_i have to be?

Analyzing SCV is better

(Dyer-Frieze'1991)

Main idea:

$$\text{SCV}(X_i) \leq \frac{\varepsilon^2/4}{t} \Rightarrow \text{SCV}(X) \lesssim \varepsilon^2/4$$

proof:

$$\text{SCV}(X) = (1 + \text{SCV}(X_1)) \dots (1 + \text{SCV}(X_t)) - 1$$

$$\text{SCV}(X) = \frac{V[X]}{E[X]^2} = \frac{E[X^2]}{E[X]^2} - 1$$

How pre

Analysis

(Dy

Main idea

$$X_1, X_2 \text{ independent} \Rightarrow E[X_1 X_2] = E[X_1]E[X_2]$$

$$X_1, X_2 \text{ independent} \Rightarrow X_1^2, X_2^2 \text{ independent}$$

$$X_1, X_2 \text{ independent} \Rightarrow$$

$$\text{SCV}(X_1 X_2) = (1 + \text{SCV}(X_1))(1 + \text{SCV}(X_2)) - 1$$

$$\text{SCV}(X_i) \leq \frac{\epsilon^2}{t} \Rightarrow \text{SCV}(X) \lesssim \epsilon^2/4$$

proof:

$$\text{SCV}(X) = (1 + \text{SCV}(X_1)) \dots (1 + \text{SCV}(X_t)) - 1$$

$$\text{SCV}(X) = \frac{V[X]}{E[X]^2} = \frac{E[X^2]}{E[X]^2} - 1$$

How precise do the X_i have to be?

Analyzing SCV is better

(Dyer-Frieze'1991)

$$X = X_1 X_2 \dots X_t$$

Main idea:

$$\text{SCV}(X_i) \leq \frac{\varepsilon^2/4}{t} \Rightarrow \text{SCV}(X) \approx \varepsilon^2/4$$

each term $O(t/\varepsilon^2)$ samples $\Rightarrow O(t^2/\varepsilon^2)$ total

Outline






6. More...

a) proof of Dyer-Frieze





b) independent sets revisited

c) warm starts








Hamiltonian



2





1







0


Hamiltonian – many possibilities



2



1



0

(hardcore lattice gas model)


**What would be a natural hamiltonian
for planar graphs?**


What would be a natural hamiltonian for planar graphs?

$H(G) = \text{number of edges}$

natural MC


pick u, v uniformly at random






natural MC

pick u, v uniformly at random





$$\pi(\mathbf{G}) \propto \lambda^{\text{number of edges}}$$

satisfies the detailed balance condition

$$\pi(\mathbf{G}) P(\mathbf{G}, \mathbf{G}') = \pi(\mathbf{G}') P(\mathbf{G}', \mathbf{G})$$

$$(\lambda = \exp(-\beta))$$

Outline

6. More...

a) proof of Dyer-Frieze

b) independent sets revisited


c) warm starts

Mixing time:

$$\tau_{\text{mix}} = \text{smallest } t \text{ such that } \|\mu_t - \pi\|_{\text{TV}} \leq 1/e$$

Relaxation time:

$$\tau_{\text{rel}} = 1/(1-\lambda_2)$$



$$\Theta(n \ln n)$$

$$\Theta(n)$$

$$\tau_{\text{rel}} \leq \tau_{\text{mix}} \leq \tau_{\text{rel}} \ln (1/\pi_{\text{min}})$$

(discrepancy may be substantially bigger for, e.g., matchings)

Estimating $\pi(S)$

Mixing time:

$$\tau_{\text{mix}} = \text{smallest } t \text{ such that } \|\mu_t - \pi\|_{\text{TV}} \leq 1/e$$

Relaxation time:


$$\tau_{\text{rel}} = 1/(1-\lambda_2)$$

$$X \sim \pi$$

$$Y = \begin{cases} 1 & \text{if } X \in S \\ 0 & \text{otherwise} \end{cases}$$

$$E[Y] = \pi(S)$$

METHOD 1



Estimating $\pi(S)$

Mixing time:

$$\tau_{\text{mix}} = \text{smallest } t \text{ such that } \|\mu_t - \pi\|_{\text{TV}} \leq 1/e$$

Relaxation time:


$$\tau_{\text{rel}} = 1/(1-\lambda_2)$$

$$X \sim \pi$$

$$Y = \begin{cases} 1 & \text{if } X \in S \\ 0 & \text{otherwise} \end{cases}$$

$$E[Y] = \pi(S)$$

METHOD 1



METHOD 2

(Gillman'98, Kahale'96, ...)



Further speed-up

Mixing time:

$$\tau_{\text{mix}} = \text{smallest } t \text{ such that } |\mu_t - \pi|_{\text{TV}} \leq 1/e$$

Relaxation time:

$$\tau_{\text{rel}} = 1/(1-\lambda_2)$$

$$|\mu_t - \pi|_{\text{TV}} \leq \exp(-t/\tau_{\text{rel}}) \text{Var}_{\pi}(\mu_0/\pi)$$

$$\left(\sum \pi(\mathbf{x}) (\mu_0(\mathbf{x})/\pi(\mathbf{x}) - 1)^2 \right)^{1/2}$$

small \Rightarrow called warm start

METHOD 2

(Gillman'98, Kahale'96, ...)



Further speed-up

sample at β can be used as a warm start for β'



cooling schedule can step from β' to β

$$|\mu_t - \pi|_{TV} \leq \exp(-t/\tau_{rel}) \text{Var}_{\pi}(\mu_0/\pi)$$

$$\left(\sum \pi(\mathbf{x}) (\mu_0(\mathbf{x})/\pi(\mathbf{x}) - 1)^2 \right)^{1/2}$$


small \Rightarrow called warm start

METHOD 2

(Gillman'98, Kahale'96, ...)



sample at β can be used as a warm start for β'



cooling schedule can step from β' to β

previous cooling schedules

$$Z(0) = A \quad H: \Omega \rightarrow \{0, \dots, n\}$$

$$\beta_0 = 0 < \beta_1 < \beta_2 < \dots < \beta_t = \infty$$

“Safe steps”

$$\beta \rightarrow \beta + 1/n$$

$$\beta \rightarrow \beta (1 + 1/\ln A)$$

$$\ln A \rightarrow \infty$$

(Bezáková, Štefankovič, Vigoda, V. Vazirani'06)


Cooling schedules of length

$$O(n \ln A)$$

$$O((\ln n)(\ln A))$$

(Bezáková, Štefankovič, Vigoda, V. Vazirani'06)

$$m = O((\ln n)(\ln A))$$




● = “well mixed” states



 = "well mixed" states

run the our cooling-schedule algorithm with METHOD 2 using "well mixed" states as starting points

METHOD 2




Output of our algorithm:

$$k = O^*((\ln A)^{1/2})$$



small augmentation (so that we can use sample from current β as a warm start at next)
still $O^*((\ln A)^{1/2})$



Use analogue of Frieze-Dyer for independent samples from vector variables with slightly dependent coordinates.

if we have sampler oracles for μ_β
then we can get adaptive schedule
of length $t=O^* \left((\ln A)^{1/2} \right)$

independent sets $O^*(n^2)$
(using Vigoda'01, Dyer-Greenhill'01)

matchings $O^*(n^2m)$
(using Jerrum, Sinclair'89)

spin systems:

Ising model $O^*(n^2)$ for $\beta < \beta_c$
(using Marinelli, Olivieri'95)

k-colorings $O^*(n^2)$ for $k > 2\Delta$
(using Jerrum'95)