

IMPORTANT: Write each problem on a separate sheet of paper. Write your name on each sheet. (Why? To speed up the grading we will use parallelization—each problem will be graded by a different TA. At the beginning of the class (on the date when the homework is due) there will be a separate pile for each problem.)

Homework 4 due: Nov. 15 (Thursday); collected before class in Wegmans 1400.

Problem sessions before Homework 4 is due:

Nov. 12 (Monday) 6:15pm - 7:15pm in Hylan 202

Nov. 13 (Tuesday) 6:15pm - 7:15pm in Goergen 109

Nov. 14 (Wednesday) 6:15pm - 7:15pm in Gavett 310

Nov. 14 (Wednesday) 7:40pm - 8:40pm in Hutchinson 473

Exam 3 is on: Nov. 29 (Thursday).

Problem sessions before Exam 3:

Nov. 26 (Monday) 6:15pm - 7:15pm in Hylan 202

Nov. 27 (Tuesday) 6:15pm - 7:15pm in Goergen 109

Nov. 27 (Wednesday) 6:15pm - 7:15pm in Gavett 310

Nov. 28 (Wednesday) 7:40pm - 8:40pm in Hutchinson 473

1 Homework - solve and turn in

4.1 (due Nov. 15) Krusty the Clown purchased a new super-efficient SUV with two engines. The first engine runs on a 2:1 mixture of gasoline and ethanol and gets 12 mpg. The second engine runs on a 1:2 mixture of gasoline and ethanol and gets 22 mpg. Krusty has 12 gallons of gasoline and 14 gallons of ethanol. How far can he get?

- (a) Write a linear program for the problem. You are allowed to have only two variables in your linear program.
- (b) Solve the linear program (you do not need to use the simplex method).
- (c) Write down the dual of your linear program from part (a).
- (d) Solve the dual program (you do not need to use the simplex method).

4.2 This problem involves three chemicals A, B, C . You have a liters of chemical A , b liters of chemical B , and c liters of chemical C (for each chemical you have a pure solution). There are n potential customers. The i -th customer wants d_i liters of an $a_i : b_i : c_i$ mixture of chemicals A, B, C and is willing to pay p_i dollars per liter of the desired mixture. You need cash. You want to get as much money as possible by creating mixtures and selling them to the potential customers. The customers will purchase your mixture even if you only partially satisfy their demand (for example, if a customer wants 3 liters of a 7 : 8 : 9 mixture and is willing to pay \$8 per liter and you supply 1.2 liters of a 7 : 8 : 9 mixture they will pay you \$9.6). Give a linear program for the problem. **Clearly describe the variables in your linear program.**

4.3 (due Nov. 15) Zofka is trying to measure the length of a race track using a car. The (digital) odometer in the car shows the number of kilometers traveled since the car was made. She drives around the track and each time she crosses the finish line she writes down the number on the odometer. She wants to infer the length of the race track from these numbers in the following sense: 1) she wants to know the minimal possible track length that is consistent with the measurements and 2) she wants to know the maximal possible track length that is consistent with the measurements.

Example input: suppose Zofka took one lap around the track. When she crossed the finish line at the beginning of the first lap the odometer showed 713. When she crossed the finish line at the end of the first lap the odometer showed 715. The minimal track length is 1 kilometer (if the car went from 713.999... km to 715.00km). The maximal track length is 3 kilometers (if the car went from 713.000km to 715.999... km).¹

The input of the problem is n —the number of laps Zofka did around the track and $n+1$ numbers $a_0, a_1, a_2, \dots, a_{n+1}$ (the values the odometer showed when she crossed the finish line). Give a linear problem that computes the minimal possible track length that is consistent with the measurements (the linear problem that computes the maximal possible track length that is consistent with the measurements is the same with min replaced by max). **The linear program should have exactly two variables.** Clearly explain the meaning of variables in your programs.

Show how your general solution works for the example input.

¹Technically we should be talking about suprema instead of maxima and infima instead of minima. Ignore this unimportant technical detail (if it worries you—then you are allowed strict inequalities in your linear program for this problem).

2 Bonus Homework - solve and turn in

4.4 (due Nov. 15)

PART A:

We are given n pairs of real numbers $(x_1, y_1), \dots, (x_n, y_n)$. We want to find real numbers a, b which minimize

$$\max_{i \in \{1, \dots, n\}} |ax_i + y_i - b|.$$

Write a linear program for this problem.

PART B:

$(x_1, y_1), \dots, (x_n, y_n)$. We want to find real numbers a, b which minimize

$$\sum_{i \in \{1, \dots, n\}} |ax_i + y_i - b|.$$

Write a linear program for this problem.

PART C:

We are given n pairs of real numbers $(x_1, y_1), \dots, (x_n, y_n)$. We want to find real numbers a, b which minimize

$$\sum_{i \in \{1, \dots, n\}} (ax_i + y_i - b)^2.$$

Find a formula for a and b (you do NOT need linear programming for this one).